

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 April 2004 (22.04.2004)

PCT

(10) International Publication Number
WO 2004/033454 A1

(51) International Patent Classification⁷: C07D 471/04,
A61K 31/437, A61P 31/22 // (C07D 471/04, 231:00,
221:00)

(74) Agents: LEVY, David, J. et al.; GlaxoSmithKline, Five
Moore Drive, PO Box 13398, Research Triangle Park, NC
27709 (US).

(21) International Application Number:
PCT/US2003/030334

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
24 September 2003 (24.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(30) Priority Data:
60/415,939 3 October 2002 (03.10.2002) US

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM CORPORATION
[US/US]; One Franklin Plaza, Philadelphia, PA 19101
(US).


Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(72) Inventors; and
(75) Inventors/Applicants (*for US only*): GUDMUNDSSON,
Kristjan, S. [IS/US]; GlaxoSmithKline, Five Moore
Drive, PO Box 13398, Research Triangle Park, NC 27709
(US). JOHNS, Brian, A. [US/US]; GlaxoSmithKline,
Five Moore Drive, PO Box 13398, Research Triangle Park,
NC 27709 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: THERAPEUTIC COMPOUNDS BASED ON PYRAZOLOPYRIDINE DERIVATIVES



(I)

(57) Abstract: The present invention provides compounds of Formula (I): wherein all variables are as defined herein, pharmaceutical compositions containing the same, processes for preparing the same and their use as pharmaceutical agents

THERAPEUTIC COMPOUNDS BASED ON PYRAZOLOPYRIDINE DERIVATIVES

5 The present invention relates to novel compounds, pharmaceutical formulations comprising these compounds, and the use of these compounds in therapy. More particularly, the present invention relates to compounds for the prophylaxis and treatment of herpes viral infections.

10 Of the DNA viruses, those of the herpes group cause the most common viral illnesses in man. The group includes herpes simplex virus types 1 and 2 (HSV), varicella zoster virus (VZV), cytomegalovirus (CMV), Epstein-Barr virus (EBV), human herpes virus type 6 (HHV-6), human herpes virus type 7 (HHV-7) and human herpes virus type 8 (HHV-8). HSV-1 and HSV-2 are some of the most common infectious agents of man. Most of
15 these viruses are able to persist in the host's neural cells; once infected, individuals are at risk of recurrent clinical manifestations of infection which can be both physically and psychologically distressing.

Herpes simplex viruses (HSV-1 and -2) are the causative agents of herpes labialis and
20 genital herpes. HSV infection is often characterised by extensive and debilitating lesions of the skin, mouth and/or genitals. Primary infections may be subclinical although tend to be more severe than infections in individuals previously exposed to the virus. Ocular infection by HSV can lead to keratitis or cataracts thereby endangering the host's sight. Infection in the new-born, in immunocompromised
25 patients or penetration of the infection into the central nervous system can prove fatal. In the US alone, 40 million individuals are infected with HSV-2, a number that is expected to increase to 60 million by 2007. Over 80% of individuals infected with HSV-2 are unaware they carry and spread the virus, and of those diagnosed less than 20% received oral therapies. The net result is that less than 5% of the infected
30 population are treated. Likewise of the 530 million individuals worldwide who carry the HSV-1 virus, 81% of the symptomatic population remain untreated. No cure

exists for HSV infection, and once infected, individuals carry the virus for life in a dormant state. Reactivation of the virus from latency occurs periodically and may be triggered by stress, environmental factors, and/or suppression of the host immune system. Currently, the use of nucleoside analogs such as valaciclovir (VALTREX®) and
5 aciclovir (ZOVIRAX®) is the standard of care for managing genital herpes virus outbreaks.

Varicella zoster virus (VZV) (also known as herpes zoster virus) is a herpes virus which causes chickenpox and shingles. Chickenpox is the primary disease produced in a host
10 without immunity, and in young children is usually a mild illness characterised by a vesicular rash and fever. Shingles or zoster is the recurrent form of the disease which occurs in adults who were previously infected with VZV. The clinical manifestations of shingles are characterised by neuralgia and a vesicular skin rash that is unilateral and dermatomal in distribution. Spread of inflammation may lead to paralysis or
15 convulsions. Coma can occur if the meninges become affected. VZV is of serious concern in patients receiving immunosuppressive drugs for transplant purposes or for treatment of malignant neoplasia and is a serious complication of AIDS patients due to their impaired immune system.


20 In common with other herpes viruses, infection with CMV leads to a lifelong association of virus and host. Congenital infection following infection of the mother during pregnancy may give rise to clinical effects such as death or gross disease (microcephaly, hepatosplenomegaly, jaundice, mental retardation), retinitis leading to blindness or, in less severe forms, failure to thrive, and susceptibility to chest and ear
25 infections. CMV infection in patients who are immunocompromised for example as a result of malignancy, treatment with immunosuppressive drugs following transplantation or infection with Human Immunodeficiency Virus, may give rise to retinitis, pneumonitis, gastrointestinal disorders and neurological diseases. CMV infection is also associated with cardiovascular diseases and conditions including
30 restenosis and atherosclerosis.

The main disease caused by EBV is acute or chronic infectious mononucleosis (glandular fever). Examples of other EBV or EBV associated diseases include lymphoproliferative disease which frequently occurs in persons with congenital or acquired cellular immune deficiency, X-linked lymphoproliferative disease which occurs namely in young boys, EBV-associated B-cell tumours, Hodgkin's disease, nasopharyngeal carcinoma, Burkitt lymphoma, non-Hodgkin's lymphoma, thymomas and oral hairy leukoplakia. EBV infections have also been found in association with a variety of epithelial-cell-derived tumours of the upper and lower respiratory tracts including the lung. EBV infection has also been associated with other diseases and conditions including chronic fatigue syndrome, multiple sclerosis and Alzheimer's disease.

HHV-6 has been shown to be a causative agent of infantum subitum in children and of kidney rejection and interstitial pneumonia in kidney and bone marrow transplant patients, respectively, and may be associated with other diseases such as multiple sclerosis. There is also evidence of repression of stem cell counts in bone marrow transplant patients. HHV-7 is of undetermined disease aetiology.

Hepatitis B virus (HBV) is a viral pathogen of world-wide major importance. The virus is aetiologically associated with primary hepatocellular carcinoma and is thought to cause 80% of the world's liver cancer. Clinical effects of infection with HBV range from headache, fever, malaise, nausea, vomiting, anorexia and abdominal pains. Replication of the virus is usually controlled by the immune response, with a course of recovery lasting weeks or months in humans, but infection may be more severe leading to persistent chronic liver disease outlined above.

U.S. Patent No. 5,498,774 and European Patent No. 0 404 190 to Mitsudera et al., relates to condensed heterocyclic compounds of the formula (I):




wherein Q is a condensed heterocyclic group having a nitrogen atom in the bridgehead which is unsubstituted or substituted, X is a hydrogen atom or a group bonded through C, O, S or N, and Y is an electron attractive group; or its salt which is useful as an agricultural chemical.

5

BRIEF SUMMARY OF THE INVENTION

The present invention provides a compound of formula (I):

10



wherein:

15

p is 0, 1, 2, 3 or 4;

each R¹ is the same or different and is independently selected from the group consisting of halo, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Ay, Het, -OR⁷, -OAY, -OR¹⁰AY, -OHet, -OR¹⁰Het, -C(O)R⁹, -C(O)AY, -C(O)Het, -CO₂R⁹, -C(O)NR⁷R⁸, -C(O)NR⁷AY, -C(O)NHR¹⁰AY, -C(O)NHR¹⁰Het, -C(S)NR⁹R¹¹, -C(NH)NR⁷R⁸, -C(NH)NR⁷AY, -S(O)_nR⁹, -S(O)_nAY, -S(O)_nHet, -S(O)₂NR⁷R⁸, -S(O)₂NR⁷AY, -NR⁷R⁸, -NR⁷AY, -NHHet, -NHR¹⁰AY, -NHR¹⁰Het, -R¹⁰cycloalkyl, -R¹⁰AY, -R¹⁰Het, -R¹⁰O-C(O)R⁹, -R¹⁰O-C(O)AY, -R¹⁰O-C(O)Het, -R¹⁰O-S(O)_nR⁹, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹, -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(O)NR⁷AY, -R¹⁰C(O)NHR¹⁰Het, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO_nR⁹, -R¹⁰SO₂NR⁹R¹¹, -R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷AY, -R¹⁰NHC(NH)NR⁹R¹¹, cyano, nitro and azido; or

25

two adjacent R¹ groups together with the atoms to which they are bonded form a C₅₋₆cycloalkyl or a 5 or 6-membered heterocyclic ring containing 1 or 2 heteroatoms;

30

each R⁷ and R⁸ are the same or different and are independently selected from the group consisting of H, alkyl, alkenyl, cycloalkyl, cycloalkenyl,

-C(O)R⁹, -CO₂R⁹, -C(O)NR⁹R¹¹, -C(S)NR⁹R¹¹, -C(NH)NR⁹R¹¹, -SO₂R¹⁰,
 -SO₂NR⁹R¹¹, -R¹⁰cycloalkyl, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹,
 -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO₂R¹⁰,
 -R¹⁰SO₂NR⁹R¹¹, -R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁹R¹¹, -R¹⁰NHCOR⁹,
 -R¹⁰NHSO₂R⁹ and -R¹⁰NHC(NH)NR⁹R¹¹;

each R⁹ and R¹¹ are the same or different and are independently selected from
 the group consisting of H, alkyl, cycloalkyl, -R¹⁰cycloalkyl, -R¹⁰OH,
 -R¹⁰(OR¹⁰)_w where w is 1-10, and -R¹⁰NR¹⁰R¹⁰;

each R¹⁰ is the same or different and is independently selected from the group
 consisting of alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkenyl;

Ay is aryl;

Het is a 5- or 6-membered heterocyclic or heteroaryl group;

R² is selected from the group consisting of halo, alkyl, alkenyl, cycloalkyl, cycloalkenyl,

Ay, Het, -OR⁷, -OAY, -OHet, -OR¹⁰Het, -S(O)_nR⁹, -S(O)_nAY, -S(O)_nNR⁷R⁸,

-S(O)_nHet, -NR⁷R⁸, -NHHet, -NHR¹⁰AY, -NHR¹⁰Het, -R¹⁰NR⁷R⁸ and -R¹⁰NR⁷AY;

n is 0, 1 or 2;

Y is N or CH;

R³ and R⁴ are the same or different and are each independently selected from the
 group consisting of H, halo, alkyl, alkenyl, cycloalkyl, Ay, Het, -OR⁷, -OAY,

-C(O)R⁷, -C(O)AY, -CO₂R⁷, -CO₂AY, -SO₂NHR⁹, -NR⁷R⁸, -NR⁷AY, -NHHet,

-NHR¹⁰Het, -R¹⁰cycloalkyl, -R¹⁰OR⁷, -R¹⁰OAY, -R¹⁰NR⁷R⁸ and -R¹⁰NR⁷AY;

R⁵ is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl,

cycloalkyl, cycloalkenyl, -OR⁷, -OAY, -OHet, -OR¹⁰AY, -OR¹⁰Het, -C(O)R⁹,

-C(O)AY, -C(O)Het, -CO₂R⁹, -C(O)NR⁷R⁸, -C(O)NR⁷AY, -C(O)NHR¹⁰Het, -CH(OR⁹)₂,

-CH(OR⁹)-R¹⁰, -CH(OR⁹)-AY, -C(S)NR⁹R¹¹, -C(NH)NR⁷R⁸, -C(NH)NR⁷AY, -S(O)_nR⁹,

-S(O)₂NR⁷R⁸, -S(O)₂NR⁷AY, -NR⁷R⁸, -NR⁷AY, -NHHet, -NHR¹⁰AY, -NHR¹⁰Het,

-R¹⁰cycloalkyl, -R¹⁰AY, -R¹⁰Het, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰C(O)AY, -R¹⁰C(O)Het,

-R¹⁰CO₂R⁹, -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(O)NR⁷AY, -R¹⁰C(O)NHR¹⁰Het, -R¹⁰CH(OR⁹)-R¹⁰;

-R¹⁰CH(OR⁹)-AY, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO_nR⁹, -R¹⁰SO₂NR⁹R¹¹,

-R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷AY, -R¹⁰NHC(NH)NR⁹R¹¹, cyano, nitro and

azido; or

wherein when Y is CH, R³ is not -NR⁷Ay;
or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

- 5 The present invention also provides a pharmaceutical composition comprising a compound of formula (I). In one embodiment, the pharmaceutical composition further comprises a pharmaceutically acceptable carrier or diluent. In one embodiment, the pharmaceutical composition further comprises an antiviral agent selected from the group consisting of aciclovir, valaciclovir and pharmaceutically
10 acceptable salts thereof.

- The present invention also provides a method for the prophylaxis or treatment of a herpes viral infection in an animal. The method comprises administering a therapeutically effective amount of a compound of formula (I) or a pharmaceutically
15 acceptable salt, solvate or physiologically functional derivative thereof. The herpes viral infection can be any of herpes simplex virus 1, herpes simplex virus 2, cytomegalovirus, Epstein Barr virus, varicella zoster virus, human herpes virus 6, human herpes virus 7, and human herpes virus 8.


- 20 The present invention also provides a method for the prophylaxis or treatment of a condition or disease associated with a herpes viral infection in an animal. The method comprises administering to the animal a therapeutically effective amount of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

25


The present invention also provides a process for preparing a compound of formula (I). The process comprises the steps of:

- a) coupling a compound of formula (II):


30



wherein X is chloro, bromo, iodo or triflate;
to a terminal alkyne of formula (III):




5 to prepare a compound of formula (IV):



10 and

b) reacting an *N*-amino pyridinium salt of formula (V):



15

wherein Z- is a counterion;

with the compound of the formula (IV) to prepare a compound of formula (I).

The present invention provides a radiolabeled compound of formula (I) or a
20 pharmaceutically acceptable salt, solvate or physiologically functional derivative
thereof. In one embodiment, the present invention provides a tritiated compound of
formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional
derivative thereof. In another aspect, the present invention provides a biotinylated
compound of formula (I) or a pharmaceutically acceptable salt, solvate or
25 physiologically functional derivative thereof.

The present invention provides a compound of formula (I) for use in therapy.

The present invention provides a compound of formula (I) for use in the prophylaxis or
30 treatment of a herpes viral infection in an animal, particularly a human.

The present invention provides a compound of formula (I) for use in the prophylaxis or treatment of a condition or disease associated with a herpes viral infection in an animal, particularly a human.

- 5 The present invention provides the use of a compound of formula (I) for the preparation of a medicament for the prophylaxis or treatment of a herpes viral infection in an animal, particularly a human.

- 10 The present invention provides the use of a compound of formula (I) for the preparation of a medicament for the treatment or prophylaxis of a condition or disease associated with a herpes viral infection in an animal, particularly a human. The present invention also provides a pharmaceutical composition comprising a compound of formula (I) for use in the prophylaxis or treatment of a herpes viral infection in an animal, particularly a human.

15

DETAILED DESCRIPTION OF THE INVENTION

- As used herein, "a compound of the invention" or "a compound of formula (I)" means
20 a compound of formula (I) or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. Similarly, with respect to isolateable intermediates such as for example, compounds of formula (IV), the phrase "a compound of formula "(number)" means a compound having that formula and pharmaceutically acceptable salts, solvates and physiologically functional derivatives
25 thereof.

- As used herein, the terms "alkyl" (and alkylene) refer to straight or branched hydrocarbon chains containing from 1 to 8 carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl,
30 isobutyl, isopropyl, and tert-butyl. Examples of "alkylene" as used herein include, but are not limited to, methylene, ethylene, propylene, butylene, and isobutylene. "Alkyl"

and "alkylene" also include substituted alkyl and substituted alkylene. The alkyl groups may be optionally substituted with one or more substituents selected from the group consisting of mercapto, nitro, cyano and halo. Perhalo alkyl, such as trifluoromethyl is one particular alkyl group.

5

As used herein, the term "cycloalkyl" refers to a non-aromatic carbocyclic ring having from 3 to 8 carbon atoms and no carbon-carbon double bonds. "Cycloalkyl" includes by way of example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. "Cycloalkyl" also includes substituted cycloalkyl. The cycloalkyl may be
10 optionally substituted on any available carbon(s) with one or more substituents selected from the group consisting of mercapto, nitro, cyano, halo and alkyl.

As used herein, the term "alkenyl" refers to straight or branched hydrocarbon chains containing from 2 to 8 carbon atoms and at least one and up to three carbon-carbon
15 double bonds. Examples of "alkenyl" as used herein include, but are not limited to ethenyl and propenyl. "Alkenyl" also includes substituted alkenyl. The alkenyl groups may optionally be substituted on any available carbon(s) with one or more substituents selected from the group consisting of mercapto, nitro, cyano, halo and alkyl.

20

As used herein, the term "cycloalkenyl" refers to a non-aromatic carbocyclic ring having from 3 to 8 carbon atoms (unless otherwise specified) and up to 3 carbon-carbon double bonds. "Cycloalkenyl" includes by way of example cyclobutenyl, cyclopentenyl and cyclohexenyl. "Cycloalkenyl" also includes substituted cycloalkenyl.
25 The cycloalkenyl may optionally be substituted on any available carbon(s) with one or more substituents selected from the group consisting of mercapto, nitro, cyano, halo and alkyl.

As used herein, the term "alkynyl" refers to straight or branched hydrocarbon chains
30 containing from 2 to 8 carbon atoms and at least one and up to three carbon-carbon triple bonds. Examples of "alkynyl" as used herein include, but are not limited to

ethynyl and propynyl. "Alkynyl" also includes substituted alkynyl. The alkynyl groups may optionally be substituted on any available carbon(s) with one or more substituents selected from the group consisting of mercapto, nitro, cyano, halo and alkyl.

5

The term "halo" or "halogen" refers to the elements fluorine, chlorine, bromine and iodine.

The term "aryl" refers to monocyclic carbocyclic groups and fused bicyclic carbocyclic groups having from 5 to 12 carbon atoms (unless otherwise specified) and having at least one aromatic ring. Examples of particular aryl groups include but are not limited to phenyl, and naphthyl. "Aryl" also includes substituted aryl. Aryl groups may optionally be substituted on any available carbon(s) with one or more substituents selected from the group consisting of halo, alkyl (including perhaloalkyl), alkenyl, cycloalkyl, cycloalkenyl, hydroxy, alkoxy, cycloalkoxy, alkylhydroxy, mercapto, amino, alkylamine, cycloalkylamine, Het, amidine, carboxy, carboxamide, sulfonamide, cyano, nitro and azido. Preferred aryl groups according to the invention include but are not limited to phenyl and substituted phenyl.

The term "heterocyclic" (or "heterocycle") refers to monocyclic saturated or unsaturated non-aromatic groups and fused bicyclic non-aromatic groups, having the specified number of members and containing 1, 2, 3 or 4 heteroatoms selected from N, O and S. Examples of particular heterocyclic groups include but are not limited to tetrahydrofuran, dihydropyran, tetrahydropyran, pyran, oxetane, thietane, 1,4-dioxane, 1,3-dioxane, 1,3-dioxalane, piperidine, piperazine, tetrahydropyrimidine, pyrrolidine, morpholine, thiomorpholine, thiazolidine, oxazolidine, tetrahydrothiopyran, tetrahydrothiophene, and the like. "Heterocyclic" also includes substituted heterocyclic. The heterocyclic groups may optionally be substituted on any available carbon(s) or heteroatom(s) with one or more substituents selected from the group consisting of halo, alkyl (including perhaloalkyl), alkenyl, cycloalkyl, cycloalkenyl, hydroxy, alkoxy, cycloalkoxy, alkylhydroxy, mercapto, amino, alkylamine,

cycloalkylamine, Het, amidine, carboxy, carboxamide, sulfonamide, cyano, nitro and azido. Preferred heterocyclic groups according to the invention include but are not limited to pyrrolidine, piperidine, morpholine, thiomorpholine and piperazine, and substituted variants thereof.

5

The term "heteroaryl" refers to aromatic monocyclic groups and aromatic fused bicyclic groups (wherein at least one ring is aromatic) having the specified number of members and containing 1, 2, 3, or 4 heteroatoms selected from N, O and S. Examples of particular heteroaryl groups include but are not limited to furan, thiophene, pyrrole, imidazole, pyrazole, triazole, tetrazole, thiazole, oxazole, isoxazole, oxadiazole, thiadiazole, isothiazole, pyridine, pyridazine, pyrazine, pyrimidine, quinoline, isoquinoline, benzofuran, benzothiophene, indole, and indazole. "Heteroaryl" also includes substituted heteroaryl. The heteroaryl groups may optionally be substituted on any available carbon(s) or heteroatom(s) with one or more substituents selected from the group consisting of halo, alkyl (including perhaloalkyl), alkenyl, cycloalkyl, cycloalkenyl, hydroxy, alkoxy, cycloalkoxy, alkylhydroxy, mercapto, amino, alkylamine, cycloalkylamine, Het, amidine, carboxy, carboxamide, sulfonamide, cyano, nitro and azido. Preferred heteroaryl groups according to the invention include but are not limited to pyridine, furan, thiophene, pyrrole, imidazole, pyrazole, and pyrimidine, and substituted variants thereof.

10

15

20


The term "members" (and variants thereof e.g., "membered") in the context of heterocyclic and heteroaryl groups refers to the total atoms, carbon and heteroatoms N, O and/or S, which form the ring. Thus, an example of a 6-membered heterocyclic ring is piperidine and an example of a 6-membered heteroaryl ring is pyridine.

25

As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s) that occur and events that do not occur.

30

The present invention provides compounds of formula formula (I):



wherein:

p is 0, 1, 2, 3 or 4;

each R¹ is the same or different and is independently selected from the group

consisting of halo, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Ay, Het, -OR⁷,
 -OAY, -OR¹⁰AY, -OHet, -OR¹⁰Het, -C(O)R⁹, -C(O)AY, -C(O)Het, -CO₂R⁹,
 -C(O)NR⁷R⁸, -C(O)NR⁷AY, -C(O)NHR¹⁰AY, -C(O)NHR¹⁰Het, -C(S)NR⁹R¹¹,
 -C(NH)NR⁷R⁸, -C(NH)NR⁷AY, -S(O)_nR⁹, -S(O)_nAY, -S(O)_nHet, -S(O)₂NR⁷R⁸,
 -S(O)₂NR⁷AY, -NR⁷R⁸, -NR⁷AY, -NHHet, -NHR¹⁰AY, -NHR¹⁰Het, -R¹⁰cycloalkyl,
 -R¹⁰AY, -R¹⁰Het, -R¹⁰O-C(O)R⁹, -R¹⁰O-C(O)AY, -R¹⁰O-C(O)Het, -R¹⁰O-S(O)_nR⁹,
 -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹, -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(O)NR⁷AY,
 -R¹⁰C(O)NHR¹⁰Het, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO_nR⁹, -R¹⁰SO₂NR⁹R¹¹,
 -R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷AY, -R¹⁰NHC(NH)NR⁹R¹¹, cyano, nitro and
 azido; or

two adjacent R¹ groups together with the atoms to which they are bonded
 form a C₅₋₆cycloalkyl or a 5 or 6-membered heterocyclic ring containing 1 or 2
 heteroatoms;

each R⁷ and R⁸ are the same or different and are independently selected from
 the group consisting of H, alkyl, alkenyl, cycloalkyl, cycloalkenyl,
 -C(O)R⁹, -CO₂R⁹, -C(O)NR⁹R¹¹, -C(S)NR⁹R¹¹, -C(NH)NR⁹R¹¹, -SO₂R¹⁰,
 -SO₂NR⁹R¹¹, -R¹⁰cycloalkyl, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹,
 -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO₂R¹⁰,
 -R¹⁰SO₂NR⁹R¹¹, -R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁹R¹¹, -R¹⁰NHCOR⁹,
 -R¹⁰NHSO₂R⁹ and -R¹⁰NHC(NH)NR⁹R¹¹;

each R⁹ and R¹¹ are the same or different and are independently selected from
 the group consisting of H, alkyl, cycloalkyl, -R¹⁰cycloalkyl, -R¹⁰OH,
 -R¹⁰(OR¹⁰)_w where w is 1-10, and -R¹⁰NR¹⁰R¹⁰;

each R^{10} is the same or different and is independently selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkenyl;

Ay is aryl;

Het is a 5- or 6-membered heterocyclic or heteroaryl group;

- 5 R^2 is selected from the group consisting of halo, alkyl, alkenyl, cycloalkyl, cycloalkenyl, Ay, Het, $-OR^7$, $-OAy$, $-OHet$, $-OR^{10}Het$, $-S(O)_nR^9$, $-S(O)_nAy$, $-S(O)_nNR^7R^8$, $-S(O)_nHet$, $-NR^7R^8$, $-NHHet$, $-NHR^{10}Ay$, $-NHR^{10}Het$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$;

n is 0, 1 or 2;

Y is N or CH;

- 10 R^3 and R^4 are the same or different and are each independently selected from the group consisting of H, halo, alkyl, alkenyl, cycloalkyl, Ay, Het, $-OR^7$, $-OAy$, $-C(O)R^7$, $-C(O)Ay$, $-CO_2R^7$, $-CO_2Ay$, $-SO_2NHR^9$, $-NR^7R^8$, $-NR^7Ay$, $-NHHet$, $-NHR^{10}Het$, $-R^{10}cycloalkyl$, $-R^{10}OR^7$, $-R^{10}OAy$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$;

- 15 R^5 is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-OR^7$, $-OAy$, $-OHet$, $-OR^{10}Ay$, $-OR^{10}Het$, $-C(O)R^9$, $-C(O)Ay$, $-C(O)Het$, $-CO_2R^9$, $-C(O)NR^7R^8$, $-C(O)NR^7Ay$, $-C(O)NHR^{10}Het$, $-CH(OR^9)_2$, $-CH(OR^9)-R^{10}$, $-CH(OR^9)-Ay$, $-C(S)NR^9R^{11}$, $-C(NH)NR^7R^8$, $-C(NH)NR^7Ay$, $-S(O)_nR^9$, $-S(O)_2NR^7R^8$, $-S(O)_2NR^7Ay$, $-NR^7R^8$, $-NR^7Ay$, $-NHHet$, $-NHR^{10}Ay$, $-NHR^{10}Het$, $-R^{10}cycloalkyl$, $-R^{10}Ay$, $-R^{10}Het$, $-R^{10}OR^9$, $-R^{10}C(O)R^9$, $-R^{10}C(O)Ay$, $-R^{10}C(O)Het$, $-R^{10}CO_2R^9$, $-R^{10}C(O)NR^9R^{11}$, $-R^{10}C(O)NR^7Ay$, $-R^{10}C(O)NHR^{10}Het$, $-R^{10}CH(OR^9)-R^{10}$, $-R^{10}CH(OR^9)-Ay$, $-R^{10}C(S)NR^9R^{11}$, $-R^{10}C(NH)NR^9R^{11}$, $-R^{10}SO_nR^9$, $-R^{10}SO_2NR^9R^{11}$, $-R^{10}SO_2NHCOR^9$, $-R^{10}NR^7R^8$, $-R^{10}NR^7Ay$, $-R^{10}NHC(NH)NR^9R^{11}$, cyano, nitro and azido; or

wherein when Y is CH, R^3 is not $-NR^7Ay$;


- 25 or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

- In one particular class of compounds of formula (I), p is 0, 1, 2 or 3. In another embodiment, p is 0, 1, or 2. In one particular embodiment, p is 0 or 1. In one particular embodiment p is 0. In another particular embodiment p is 1. In one embodiment, p is 2 and optionally, two adjacent R^1 groups together with the atoms
- 30

which they are bonded, form a C₅₋₆ cycloalkyl or 5- or 6-membered heterocyclic group. The phrase "two adjacent R¹ groups" refers to two R¹ groups, each bonded to adjacent carbon atoms on the pyrazolopyridine ring. In the embodiment where two adjacent R¹ groups together with the atoms to which they are bonded form a cycloalkyl or
5 heterocyclic group, p is typically 2, 3 or 4; more typically 2.

R¹ may be in the 4, 5, 6 or 7 position. In one embodiment, p is 1 or more and R¹ is in the C-7 position. In one embodiment, p is 1 or more and R¹ is in the C-5 position. In one embodiment p is 2 or more and one R¹ is in the C-5 position and one R¹ is in the
10 C-7 position.

In the embodiments where two adjacent R¹ groups together with the atoms to which they are bonded form a C₅₋₆cycloalkyl or 5- or 6-membered heterocyclic group having 1 or 2 heteroatoms, each R¹ group may be the same or different and is typically
15 selected from the group consisting of alkyl, alkenyl, -OR⁷, -S(O)_nR⁹ and -NR⁷R⁸. For example, in one embodiment two adjacent R¹ groups are -OR⁷ and together with the atoms to which they are bonded, they form a heterocyclic group such as:




20

In another embodiment, two adjacent R¹ groups are alkyl and together with the atoms to which they are bonded, they form a cycloalkyl group such as:



25

In another embodiment two adjacent R¹ groups are defined as -OR⁷ and -NR⁷R⁸, respectively and together with the atoms to which they are bonded, they form a heterocyclic group such as:



30

From these examples, additional embodiments can be readily ascertained by those skilled in the art.

In one embodiment, two adjacent R¹ groups together with the atoms to which they are bonded do not form a C₃-cycloalkyl or a 5 or 6-membered heterocyclic ring containing 1 or 2 heteroatoms.

In one embodiment, each R¹ is the same or different and is independently selected from the group consisting of halo, alkyl, cycloalkyl, Ay, Het, -OR⁷, -C(O)R⁹, -C(O)Het, -CO₂R⁹, -C(O)NR⁷R⁸, -C(O)NR⁷Ay, -C(O)NHR¹⁰Het, -S(O)_nR⁹, -S(O)₂NR⁷R⁸, -S(O)₂NR⁷Ay, -NR⁷R⁸, -NR⁷Ay, -NHHet, -NHR¹⁰Ay, -NHR¹⁰Het, -R¹⁰cycloalkyl, -R¹⁰Het, -R¹⁰OR⁹, -R¹⁰C(O)NR⁷Ay, -R¹⁰SO₂NHCOR⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷Ay, cyano, nitro and azido, or any subset thereof. More particularly, each R¹ is the same or different and is independently selected from the group consisting of halo, alkyl, Ay, Het, -OR⁷, -S(O)_nR⁹, -NR⁷R⁸, -NR⁷Ay, -NHHet, -NHR¹⁰Ay and -NHR¹⁰Het. In one particular embodiment, each R¹ is the same or different and is independently selected from the group consisting of halo, alkyl, Ay, Het, -OR⁷, -S(O)_nR⁹, -NR⁷R⁸, -NR⁷Ay, -NHHet and -NHR¹⁰Het, or any subset thereof. Particular compounds of formula (I) are defined wherein each R¹ is the same or different and is independently selected from the group consisting of halo, Ay, Het, -NR⁷R⁸ and -NR⁷Ay, or any subset thereof. In one embodiment, each R¹ is the same or different and is -NR⁷R⁸.

More specific examples of embodiments of the present invention are defined wherein each R¹ is the same or different and is independently selected from the group consisting of halo, phenyl, Het (and substituted Het), -O-alkyl, -S-alkyl, -NH₂, -NHalkyl, -NHcycloalkyl, -N(alkyl)(alkyl), -NHalkyl-OH, -NHalkyl-O-alkyl, -NHAy, and -NH-alkyl-Het, or any subset thereof. Specific examples of some particular R¹ groups are selected from the group consisting of Br, Cl, phenyl, -NH-methyl, -NH-butyl, -N(CH₃)₂, -NH-cyclopentyl, -NH-cyclopropyl, -NH-isopropyl, -NH-phenyl, -NH(CH₂)₂OCH₃, pyrrolidine, and morpholine, or any subset thereof.

In one embodiment, R^2 is selected from the group consisting of halo, alkenyl, cycloalkyl, cycloalkenyl, Ay, Het, $-OR^7$, $-OAY$, $-OHet$, $-OR^{10}Het$, $-S(O)_nR^9$, $-NR^7R^8$, $-NHHet$, $-NHR^{10}Het$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$, or any subset thereof. In particular, R^2 is selected from the group consisting of halo, Het, $-S(O)_nR^9$, $-NR^7R^8$, $-NHHet$ and $-NHR^{10}Het$, or any subset thereof. More particularly, R^2 is selected from the group consisting of $-NR^7R^8$, Het, $-NHR^{10}Het$ and $-NHHet$, or any subset thereof. In one particular embodiment, compounds of formula (I) are defined where R^2 is $-NR^7R^8$.

Specific examples of embodiments of the present invention are defined wherein R^2 is selected from the group consisting of $-NH_2$, $-NH$ -alkyl, $-NH$ -cycloalkyl, Het, $-NHHet$ and $-NH$ -alkyl-Het, or any subset thereof. Particular embodiments include those compounds of formula (I) wherein R^2 is $-NH_2$, $-NH$ -propyl, $-NH$ -isopropyl, $-NH$ -cyclopropyl, $-NH$ -butyl, $-NH$ -isobutyl, $-NH$ -cyclobutyl, $-NH$ -cyclopentyl, $-NH$ -cyclohexyl, $-NH(CH_2)_2OCH_3$, pyrrolidine (e.g., pyrrolidine bonded through N), and morpholine (e.g., morpholine bonded through N), or any subset thereof.

In one embodiment, R^7 and R^8 are each the same or different and are independently selected from the group consisting of H, alkyl, cycloalkyl, $-C(O)R^9$, $-R^{10}$ -cycloalkyl, $-R^{10}OR^9$, $-R^{10}CO_2R^9$ and $-R^{10}NR^9R^{11}$, or any subset thereof. More particularly, R^7 and R^8 are each the same or different and are independently selected from the group consisting of H, alkyl, cycloalkyl, and $-R^{10}$ -cycloalkyl, or any subset thereof. In one embodiment, R^7 and R^8 are each the same or different and are independently selected from the group consisting of H, alkyl and cycloalkyl.

The group $-R^{10}(OR^{10})_w$ in the definition of R^9 and R^{11} refers to a PEG-like chain such as alkyl-O-alkyl-O-alkyl. In one embodiment, R^9 and R^{11} are each the same or different and are independently selected from the group consisting of H, alkyl, cycloalkyl, and $-R^{10}$ -cycloalkyl, or any subset thereof. More particularly, R^9 and R^{11} are each the same or different and are independently selected from the group consisting of H and alkyl.

In one embodiment, R^{10} is alkyl or cycloalkyl; more particularly alkyl.

In one class of compounds of formula (I), Y is CH. In another class of compounds of formula (I), Y is N.

5

In one embodiment, R^3 is selected from the group consisting of H, halo, alkyl, Ay, $-OR^7$, $-CO_2R^7$, $-NR^7R^8$, $-R^{10}OR^7$ and $-R^{10}NR^7R^8$, or any subset thereof. More particularly, R^3 is selected from the group consisting of H, halo, alkyl, $-OR^7$, and $-NR^7R^8$, or any subset thereof. In one embodiment, R^3 is H or alkyl. In one embodiment R^3 is H.

10

In one embodiment, R^4 is selected from the group consisting of H, halo, alkyl, Ay, $-OR^7$, $-CO_2R^7$, $-NR^7R^8$, $-R^{10}OR^7$ and $-R^{10}NR^7R^8$, or any subset thereof. More preferably, R^4 is selected from the group consisting of H, halo, alkyl, $-OR^7$, and $-NR^7R^8$, or any subset thereof. In one embodiment, R^4 is H or alkyl. In one embodiment R^4 is H.

15

In one embodiment, R^5 is selected from the group consisting of halo, alkyl, cycloalkyl, $-OR^7$, $-C(O)R^9$, $-C(O)Ay$, $-C(O)Het$, $-CH(OR^9)-R^{10}$, $-CH(OR^9)-Ay$, $-S(O)_nR^9$, $-S(O)_2NR^7R^8$, $-NR^7R^8$, $-NR^7Ay$, $-R^{10}cycloalkyl$, $-R^{10}Ay$, $-R^{10}Het$, $-R^{10}OR^9$, $-R^{10}C(O)R^9$, $-R^{10}SO_2NR^9R^{11}$ and $-R^{10}NR^7R^8$, or any subset thereof. More particularly, R^5 is selected from the group consisting of alkyl, cycloalkyl, $-C(O)R^9$, $-C(O)Ay$, $-C(O)Het$, $-CH(OR^9)-Ay$, $-NR^7R^8$, $-R^{10}cycloalkyl$, $-R^{10}Ay$, $-R^{10}Het$, $-R^{10}OR^9$ and $-R^{10}NR^7R^8$, or any subset thereof. In one embodiment, R^5 is selected from the group consisting of alkyl, $-C(O)Ay$, $-CH(OR^9)-Ay$, $-R^{10}cycloalkyl$, $-R^{10}Ay$, $-R^{10}OR^9$ and $-R^{10}NR^7R^8$, or any subset thereof.

20

Specific examples of compounds of formula (I) are defined wherein R^5 is selected from the group consisting of alkyl (e.g., methyl, ethyl, propyl, isopropyl and isobutyl), cyclopropyl, alkyl-OH, alkyl-O-alkyl, alkyl-O-alkyl-O-alkyl, C(O)-phenyl, C(O)-benzyl, benzyl, -alkyl-Het (e.g., $-CH_2$ -morpholino and $-CH_2$ -pyrrolidiny), $-N(H)alkyl$, $-N(alkyl)_2$, $-alkyl-N(H)alkyl$, and $-alkyl-N(alkyl)_2$, or any subset thereof.

30

It is to be understood that the present invention includes all combinations and subsets of the particular and preferred groups defined hereinabove.

Specific examples of compounds of formula (I) include but are not limited to:

- 5 3-(2-Fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine;
 N-Cyclopentyl-4-(2-propylpyrazolo[1,5-*a*]pyridin-3-yl)pyridin-2-amine;
 7-Chloro-3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine;
 N-Cyclopentyl-3-[2-(cyclopentylamino)pyridin-4-yl]-2-propylpyrazolo[1,5-*a*]pyridin-
 7-amine;
- 10 2-Isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;
 2-Isobutyl-3-[2-(methylsulfinyl)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;
 N-Cyclopentyl-4-(2-isobutylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine;
 N-Cyclopentyl-4-[2-isobutyl-7-(methylthio)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-
 amine;
- 15 *N*-Cyclopentyl-4-[2-isobutyl-7-(methylsulfinyl)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-
 2-amine;
 N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isobutylpyrazolo[1,5-
 a]pyridin-7-amine;
 2-(Diethoxymethyl)-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;
- 20 3-[2-(Methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine-2-carbaldehyde;
 {3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol;
 {3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-
 yl}(phenyl)methanol;
 {3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-
 yl}(phenyl)methanone;
- 25 {7-(Cyclopentylamino)-3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-
 2-yl}(phenyl)methanone;
 4-(2-Benzylpyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine;
 4-(2-Benzyl-7-chloropyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine;
- 30 *N*-{4-[2-Benzyl-7-(cyclopentylamino)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinyl}-*N*-
 cyclopentylamine;

- N*-Cyclopentyl-4-[2-(methoxymethyl)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinamine;
N-Cyclopentyl-4-[2-(methoxymethyl)-7-(methylsulfanyl)pyrazolo[1,5-*a*]pyridin-3-yl]-
2-pyrimidinamine;
- 5 *N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-
(methoxymethyl)pyrazolo[1,5-*a*]pyridin-7-amine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(1-pyrrolidinyl)propyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
- N*-[3-[2-(Methylsulfanyl)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl]methyl-2-
propanamine;
- 10 *N*-Cyclopentyl-4-{2-[(isopropylamino)methyl]pyrazolo[1,5-*a*]pyridin-3-yl}-2-
pyrimidinamine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(isopropylamino)methyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
- 15 4-{7-Chloro-2-[3-(isopropylamino)propyl]pyrazolo[1,5-*a*]pyridin-3-yl}-*N*-
cyclopentyl-2-pyrimidinamine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(isopropylamino)propyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
- 20 4-{7-Chloro-2-[(2-methoxyethoxy)methyl]pyrazolo[1,5-*a*]pyridin-3-yl}-*N*-
cyclopentyl-2-pyrimidinamine;
3-[2-(Cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)methyl]-*N*-(2-
methoxyethyl)pyrazolo[1,5-*a*]pyridin-7-amine;
- N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)-
methyl]pyrazolo[1,5-*a*]pyridin-7-amine;
- N*-Cyclopentyl-4-(2-isopropylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine;
- 25 *N*-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isopropylpyrazolo[1,5-
a]pyridin-7-amine;
- 2-Cyclopropyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;
N-Cyclopentyl-4-(2-cyclopropylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine; and
N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-cyclopropylpyrazolo[1,5-
a]pyridin-7-amine;
- 30

and pharmaceutically acceptable salts, solvates and physiologically functional derivatives thereof.

It will be appreciated by those skilled in the art that the compounds of the present invention may also be utilized in the form of a pharmaceutically acceptable salt solvate or physiologically functional derivative thereof. The pharmaceutically acceptable salts of the compounds of formula (I) include conventional salts formed from pharmaceutically acceptable inorganic or organic acids or bases as well as quaternary ammonium salts. More specific examples of suitable acid salts include hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, perchloric, fumaric, acetic, propionic, succinic, glycolic, formic, lactic, maleic, tartaric, citric, palmoic, malonic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, fumaric, toluenesulfonic, methanesulfonic (mesylate), naphthalene-2-sulfonic, benzenesulfonic hydroxynaphthoic, hydroiodic, malic, steroic, tannic and the like. In one embodiment, the compounds of formula (I) are in the form of the mesylate salt. Other acids such as oxalic, while not in themselves pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable salts. More specific examples of suitable basic salts include sodium, lithium, potassium, magnesium, aluminium, calcium, zinc, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, N-methylglucamine and procaine salts.

The term "solvate" as used herein refers to a complex of variable stoichiometry formed by a solute (a compound of formula (I)) and a solvent. Solvents, by way of example, include water, methanol, ethanol, or acetic acid.

The term "physiologically functional derivative" as used herein refers to any pharmaceutically acceptable derivative of a compound of the present invention, for example, an ester or an amide of a compound of formula (I), which upon administration to an animal, particularly a mammal, such as a human, is capable of providing (directly or indirectly) a compound of the present invention or an active

metabolite thereof. See, for example, Burger's Medicinal Chemistry And Drug Discovery, 5th Edition, Vol 1: Principles And Practice.

Processes for preparing pharmaceutically acceptable salts, solvates and physiologically functional derivatives of the compounds of formula (I) are conventional in the art. See, e.g., Burger's Medicinal Chemistry And Drug Discovery 5th Edition, Vol 1: Principles And Practice.

As will be apparent to those skilled in the art, in the processes described below for the preparation of compounds of formula (I), certain intermediates, may be in the form of pharmaceutically acceptable salts, solvates or physiologically functional derivatives of the compound. Those terms as applied to any intermediate employed in the process of preparing compounds of formula (I) have the same meanings as noted above with respect to compounds of formula (I). Processes for preparing pharmaceutically acceptable salts, solvates and physiologically functional derivatives of such intermediates are known in the art and are analogous to the processes for preparing pharmaceutically acceptable salts, solvates and physiologically functional derivatives of the compounds of formula (I).

Certain compounds of formula (I) and intermediates used in the processes of preparing compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit *cis-trans* isomerism). The individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centers are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

The present invention further provides compounds of formula (I) for use in medical therapy, e.g. in the treatment or prophylaxis, including suppression of recurrence of symptoms, of a viral disease in an animal, e.g. a mammal such as a human. The compounds of formula (I) are especially useful for the treatment or prophylaxis of viral diseases such as herpes viral infections. Herpes viral infections include, for example, herpes simplex virus 1 (HSV-1), herpes simplex virus 2 (HSV-2), cytomegalovirus (CMV), Epstein Barr virus (EBV), varicella zoster virus (VZV), human herpes virus 6 (HHV-6), human herpes virus 7 (HHV-7), and human herpes virus 8 (HHV-8). Thus, the compounds of the invention are also useful in the treatment or prophylaxis of the symptoms or effects of herpes virus infections.

The compounds of the invention are useful in the treatment or prophylaxis of conditions or diseases associated with herpes virus infections, particularly conditions or diseases associated with latent herpes virus infections in an animal, e.g., a mammal such as a human. By conditions or diseases associated with herpes viral infections is meant a condition or disease, excluding the viral infection per se, which results from the presence of the viral infection, such as chronic fatigue syndrome which is associated with EBV infection; and multiple sclerosis which has been associated with herpes viral infections such as EBV and HHV-6. Further examples of such conditions or diseases are described in the background section above.

In addition to those conditions and diseases, the compounds of the present invention may also be used for the treatment or prophylaxis of cardiovascular diseases and conditions associated with herpes virus infections, in particular atherosclerosis, coronary artery disease and restenosis and specifically restenosis following angioplasty (RFA). Restenosis is the narrowing of the blood vessels which can occur after injury to the vessel wall, for example injury caused by balloon angioplasty or other surgical and/or diagnostic techniques, and is characterized by excessive proliferation of smooth muscle cells in the walls of the blood vessel treated. It is thought that in many patients suffering from restenosis following angioplasty (RFA), viral infection, particularly by CMV and/or HHV-6 plays a pivotal role in the proliferation of the

smooth muscle cells in the coronary vessel. Restenosis can occur following a number of surgical and/or diagnostic techniques, for example, transplant surgery, vein grafting, coronary by-pass grafting and, most commonly following angioplasty.

- 5 There is evidence from work done both *in vitro* and *in vivo*, indicating that restenosis is a multifactorial process. Several cytokines and growth factors, acting in concert, stimulate the migration and proliferation of vascular smooth muscle cells (SMC) and production of extracellular matrix material, which accumulate to occlude the blood vessel. In addition growth suppressors act to inhibit the proliferation of SMC's and
10 production of extracellular matrix material.

In addition, compounds of formula (I) may be useful in the treatment or prophylaxis of conditions or diseases associated with hepatitis B or hepatitis C viruses, human papilloma virus (HPV) and HIV.

15

The present invention also provides a method for the treatment or prophylaxis of a viral infection in an animal such as a mammal (e.g., a human), particularly a herpes viral infection, which method comprises administering to the animal a therapeutically effective amount of the compound of formula (I).

20

As used herein, the term "prophylaxis" refers to the prevention of infection, the prevention of occurrence of symptoms in an infected subject, the prevention of recurrence of symptoms in an infected subject, or a decrease in severity or frequency of symptoms of viral infection, condition or disease in the subject.

25

As used herein, the term "treatment" refers to the partial or total elimination of symptoms or decrease in severity of symptoms of viral infection, condition or disease in the subject, or the elimination or decrease of viral presence in the subject.

30

As used herein, the term "therapeutically effective amount" means an amount of a compound of formula (I) which is sufficient, in the subject to which it is administered,

to treat or prevent the stated disease, condition or infection. For example, a therapeutically effective amount of a compound of formula (I) for the treatment of a herpes virus infection is an amount sufficient to treat the herpes virus infection in the subject.

5

The present invention also provides a method for the treatment or prophylaxis of a condition or disease associated with herpes viral infections in an animal such as a mammal (e.g., a human), which comprises administering to the animal a therapeutically effective amount of the compound of formula (I). In one embodiment,
10 the present invention provides a method for the treatment or prophylaxis of chronic fatigue syndrome and multiple sclerosis in an animal such as a mammal (e.g., a human), which comprises administering to the animal a therapeutically effective amount of a compound of formula (I). The foregoing method is particularly useful for the treatment or prophylaxis of chronic fatigue syndrome and multiple sclerosis
15 associated with latent infection with a herpes virus.

In another embodiment, the present invention provides a method for the treatment or prophylaxis of a cardiovascular condition such as atherosclerosis, coronary artery disease or restenosis (particularly restenosis following surgery such as angioplasty),
20 which comprises administering to the animal a therapeutically effective antiviral amount of the compound of formula (I).

The present invention further provides a method for the treatment or prophylaxis of hepatitis B or hepatitis C viruses in an animal such as a mammal (e.g., a human), which
25 comprises administering to the animal a therapeutically effective amount of the compound of formula (I).

The present invention further provides a method for the treatment or prophylaxis of human papilloma virus in an animal such as a mammal (e.g., a human), which
30 comprises administering to the animal a therapeutically effective amount of the compound of formula (I).

The present invention further provides a method for the treatment or prophylaxis of HIV in an animal such as a mammal (e.g., a human), which comprises administering to the animal a therapeutically effective amount of the compound of formula (I).

- 5 The present invention also provides the use of the compound of formula (I) in the preparation of a medicament for the treatment or prophylaxis of a viral infection in an animal such as a mammal (e.g., a human), particularly a herpes viral infection; the use of the compound of formula (I) in the preparation of a medicament for the treatment of a condition or disease associated with a herpes viral infection; and the
- 10 use of the compound of formula (I) in the preparation of a medicament for the treatment or prophylaxis of hepatitis B or hepatitis C viruses, human papilloma virus and HIV. In particular, the present invention also provides the use of a compound of formula (I) in the preparation of a medicament for the treatment or prophylaxis of chronic fatigue syndrome or multiple sclerosis. In one embodiment, the present
- 15 invention provides the use of a compound of formula (I) in the preparation of a medicament for the treatment or prophylaxis of cardiovascular disease, such as restenosis and atherosclerosis.

- The compounds of formula (I) are conveniently administered in the form of
- 20 pharmaceutical compositions. Such compositions may conveniently be presented for use in conventional manner in admixture with one or more physiologically acceptable carriers or diluents.

- While it is possible that compounds of the present invention may be therapeutically
- 25 administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical composition. The pharmaceutical composition may comprise a pharmaceutically acceptable carrier or diluent. The carrier(s) or diluent(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Accordingly, the present invention further provides for a pharmaceutical formulation or composition comprising a compound of formula (I) with one or more pharmaceutically acceptable carriers therefore and, optionally, other therapeutic and/or prophylactic ingredients.

5

The formulations include those suitable for oral, parenteral (including subcutaneous e.g. by injection or by depot tablet, intradermal, intrathecal, intramuscular e.g. by depot and intravenous), rectal and topical (including dermal, buccal and sublingual) administration although the most suitable route may depend upon for example the condition, age, and disorder of the recipient as well as the viral infection or disease being treated. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association the compound(s) ("active ingredient") with the carrier which constitutes one or more accessory ingredients. In the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations suitable for oral administration may be presented as discrete units such as capsules (including soft-gel capsules), cachets or tablets (e.g. chewable tablets in particular for paediatric administration) each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with other conventional excipients such as binding agents, (for example, syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch or

30

polyvinylpyrrolidone), fillers (for example, lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate or sorbitol), lubricants (for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica), disintegrants (for example, potato starch or sodium starch glycollate) or wetting agents, such as sodium lauryl sulfate. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. The tablets may be coated according to methods well-known in the art.

10

Alternatively, the compounds of the present invention may be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, for example. Moreover, formulations containing these compounds may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminum stearate gel or hydrogenated edible fats; emulsifying agents such as lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may include edible oils) such as almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol; and preservatives such as methyl or propyl p-hydroxybenzoates or sorbic acid. Such preparations may also be formulated as suppositories, e.g., containing conventional suppository bases such as cocoa butter or other glycerides. Liquid preparations may also be formulated as soft-gel capsules for oral administration, e.g., containing conventional soft-gel excipients such as polyethylene glycol.

25

Formulations for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

30

The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of a sterile liquid carrier, for example, water-for-injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter, hard fat or polyethylene glycol.

- 10 Formulations suitable for topical (e.g., dermal) or intranasal application include ointments, creams, lotions, pastes, gels, sprays, aerosols and oils. Suitable carriers for such formulations include petroleum jelly, lanolin, polyethyleneglycols, alcohols, and combinations thereof.
- 15 Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured base such as sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a base such as gelatin and glycerin or sucrose and acacia.
- 20 The compounds may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

In addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.

It be appreciated that the amount of a compound of the invention required for use in treatment will vary with the nature of the condition being treated and the age and the condition of the patient and will be ultimately at the discretion of the attendant physician or veterinarian. In general, however, doses employed for adult human
5 treatment will typically be in the range of 0.02-5000 mg per day, preferably 100-1500 mg per day. The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example as two, three, four or more sub-doses per day. The formulations according to the invention may contain
10 between 0.1-99% of the active ingredient, conveniently from 30-95% for tablets and capsules and 3-50% for liquid preparations.

The compound of formula (I) for use in the instant invention may be used in combination with other therapeutic agents for example, non-nucleotide reverse transcriptase inhibitors, nucleoside reverse transcriptase inhibitors, protease inhibitors
15 and/or other antiviral agents. The invention thus provides in a further aspect the use of a combination comprising a compound of formula (I) with a further therapeutic agent in the treatment of viral infections. Particular antiviral agents which may be combined with the compounds of the present invention include aciclovir, valaciclovir, famcyclovir, ganciclovir, docosanol, miribavir, amprenavir, lamivudine, zidovudine, and
20 abacavir. Preferred antiviral agents for combining with the compounds of the present invention include aciclovir and valaciclovir. Thus the present invention provides in a further aspect, a combination comprising a compound of formula (I) and an antiviral agent selected from the group consisting of aciclovir or valaciclovir; the use of such combination in the treatment of viral infections and the preparation of a medicament
25 for the treatment of viral infections, and a method of treating viral infections comprising administering a compound of formula (I) and an antiviral agent selected from the group consisting of aciclovir and valaciclovir.


When the compounds of formula (I) are used in combination with other therapeutic
30 agents, the compounds may be administered either sequentially or simultaneously by any convenient route.

The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above optionally together with a pharmaceutically acceptable carrier or diluent comprise a further aspect of the invention. The individual
5 components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations..

When combined in the same formulation it will be appreciated that the two compounds must be stable and compatible with each other and the other components
10 of the formulation and may be formulated for administration. When formulated separately they may be provided in any convenient formulation, in such a manner as are known for such compounds in the art.

When a compound of formula (I) is used in combination with a second therapeutic
15 agent active against the viral infection, the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily appreciated by those skilled in the art.

Compounds of formula (I) may be conveniently prepared by a process outlined in
20 Scheme 1 below.




wherein X is chloro, bromo, iodo or triflate, and Z⁻ is a counterion (e.g., a halide,
30 sulfonate, etc.) and all other variables are as defined above.

Generally, the process for preparing a compound of formula (I) comprises the steps of:

- a) coupling a compound of formula (II) to a terminal alkyne of formula (III) to prepare a compound of formula (IV); and
- b) reacting an *N*-amino pyridinium salt of formula (V) with the compound of the formula (IV) to prepare a compound of formula (I).

More specifically, a compound of formula (I) can be prepared by reacting a compound of formula (IV) with an *N*-amino pyridinium of formula (V).



15 wherein all variables are as defined above:


This method can be readily carried out by mixing a compound of the formula (IV) in near equimolar amounts with a compound of formula (V) in an inert solvent in the presence of a base optionally with cooling to effect the cycloaddition reaction.

Suitable solvents include but are not limited to acetonitrile, dimethyl sulfoxide,

20 tetrahydrofuran, *N,N*-dimethylformamide, dioxane, 1-methyl-2-pyrrolidinone, and the lower alcohols such as methanol and ethanol. Typically the base is an amine base such as triethylamine and the like or an inorganic base such as sodium hydroxide, sodium bicarbonate, potassium carbonate, or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) and the like. It is currently believed that the initial cycloaddition reaction produces a


25 dihydro intermediate which is not generally observed, and which undergoes air oxidation to the aromatic compound of formula (I).

Compounds of formula (V) are aminated pyridine derivatives and are either commercially available or can be conveniently prepared by reacting a suitable pyridine with an aminating reagent such as O-(mesitylsulfonyl)hydroxylamine and the like.



wherein all variables are as defined above.

- 10 A compound of the formula (IV) can be prepared by coupling (e.g., cross-coupling) a compound of formula (II) with a terminal alkyne of formula (III).




wherein all variables are as defined above.

- This reaction is typically performed in an inert solvent in the presence of catalytic amounts of both a palladium (0) source and a copper (I) source along with a stoichiometric amount of base. Palladium catalysts include but are not limited to tetrakis(triphenylphosphine) palladium (0) and dichlorobis(triphenylphosphine) palladium (II). Copper (I) iodide is one source of copper co-catalyst. Typically the base is a secondary or tertiary amine such as diisopropylamine, triethylamine or diisopropylethylamine. Typical solvents are tetrahydrofuran, *N,N*-dimethylformamide, dioxane, and 1-methyl-2-pyrrolidinone and the like. This method is known in the art as a Sonogashira coupling. Accordingly, other catalysts, bases and solvents which can be employed for Sonogashira couplings may similarly be adapted for use in the preparation of the compounds of the present invention.

Compounds of the formula (II) are either commercially available or can be prepared according to methods known to one skilled in the art or by procedures described in the literature.

- 5 In addition to the foregoing process for preparing compounds of formula (I), the present invention also provides certain intermediate compounds for use in the preparation of such compounds of formula (I) according to the foregoing process. Such intermediates are described in connection with Scheme 1 above.
- 10 Each of the foregoing processes may further comprise the step of converting the compound of formula (I) to a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, using techniques well known to those skilled in the art.
- 15 As will be apparent to those skilled in the art, a compound of formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof may be converted to another compound of formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof using techniques well known in the art. For example, one method of converting a
- 20 compound of formula (I) to another compound of formula (I) comprises converting a compound of formula (I-A) to a compound of formula (I-B) via a deprotonation electrophile addition reaction.



wherein q is 0, 1, or 2;


R¹ is as defined above and is located at the C-4, C-5 and/or C-6 position; and

- 30 R^{1a} is selected from halo, alkyl, -C(O)R⁹, -C(O)Ay, -C(O)Het, -CO₂R⁹, -C(O)NR⁷R⁸, -C(O)NR⁷Ay, -C(O)NHR¹⁰Het, -S(O)₂NR⁷R⁸, -S(O)₂NR⁷Ay and -S(O)_nR⁹; and

all other variables are as defined above.

More specifically a compound of formula (I-B) can be prepared by treating a compound of formula (I-A) with a base in an inert solvent followed by the addition of an electrophilic quenching agent. Suitable bases include by way of example lithium amide bases such as lithium diisopropyl amide, and alkyllithium bases such as butyllithium. Suitable solvents include but are not limited to ether, tetrahydrofuran and the like. Electrophilic quenching reagents include but are not limited to carbon tetrachloride, hexachloroethane ($R^{1a}=Cl$), *N*-bromosuccinimide ($R^{1a}=Br$), and dimethyldisulfide ($R^{1a}=-SMe$).

Another particular process for converting a compound of formula (I) to another compound of formula (I) involves the deprotonation/electrophilic quench described above wherein the electrophile is *B*-OCH₃-9-borabicyclo[3.3.1]nonane (*B*-OMe-9-BBN). A compound of formula (I-A) thus forms upon lithiation of C7 and quenching with *B*-OMe-9-BBN an intermediate arylborinate of formula (VII) *in situ*. The aryl borinate is then able to be cross-coupled with an aryl halide in a palladium catalyzed Suzuki coupling optionally in the presence of a base to provide a compound of formula (I-C).



q is 0, 1, or 2;


R¹ is as defined above and is located at the C-4, C-5 and/or C-6 position;

R^{1b} is selected from Ay, Het, alkenyl, or cycloalkenyl; and

all other variables are as defined above.

5

As a further example, a compound of formula (I-D) can be converted to a compound of formula (I-F) by a) oxidizing the compound of formula (I-D) to prepare a compound of formula (I-E) and then b) optionally reacting a compound of formula (I-E) with an oxygen or amine nucleophile of formula R², wherein R² is selected from the group consisting of -NR⁷R⁸, -OR⁷, -OAr, Het attached through N, -NHHet, -NHR¹⁰Het, -OHet and -OR¹⁰Het to produce a compound of formula (I-F) wherein R² is selected from the group consisting of -NR⁷R⁸, -OR⁷, -OAr, Het attached through N, -NHHet, -NHR¹⁰Het, -OHet and -OR¹⁰Het.



wherein n' is 1 or 2 and all other variables are as defined above.

20

More specifically, a compound of formula (I-F) can be prepared by reacting a compound of formula (I-E) (i.e., a compound of formula (I) wherein R² is S(O)_nR⁹ where n' is 1 or 2) with an oxygen or amine nucleophile of formula R², wherein R² is selected from the group consisting of -NR⁷R⁸, -OR⁷, -OAr, Het linked through N, -NHHet, -NHR¹⁰Het, -OHet, and -OR¹⁰Het. The reaction may be carried out neat or in a suitable solvent and may be heated to 50-150°C. Typically the solvent is a lower alcohol such as methanol, ethanol, isopropanol or the like or a solvent such as *N,N*-dimethylformamide or tetrahydrofuran, or the like. Optionally a base may be used to facilitate the reaction. Typically the base can be potassium carbonate, or an amine base such as triethylamine.


25

30

A compound of formula (I-E) may be conveniently prepared by reacting a compound of formula (I-D) (i.e., a compound of formula (I) wherein R^2 is $S(O)_nR^9$ where n is 0) with an oxidizing agent in an inert solvent, optionally in the presence of an acid or base. Typically the oxidizing agent is a peracid such as *m*-chloroperbenzoic acid or the like, optionally with a base such as sodium bicarbonate. If additional amines are present in the molecule, it may be desirable to perform the oxidation under acidic conditions to protect the amine from overoxidation. By way of example some specific examples of suitable acids include acetic acid, hydrochloric acid and the like. Careful monitoring of the stoichiometry between the oxidizing agent and the substrate allows the product distribution between sulfoxide ($n=1$), and sulfone ($n=2$) to be controlled. Suitable solvents include but are not limited to, dichloromethane, chloroform and the like.

A compound of formula (I-D) can be prepared by the method described above wherein $R^2=SR^9$ from the reaction of compounds selected from the group consisting of compounds of formula (IV), compounds of formula (V) prepared from compounds of formula (II) and formula (III) (i.e., the compound of formula (II) wherein R^2 is SR^9). The requisite compound of formula (II) can be obtained from commercial sources or prepared by methods known to one skilled in the art.


Another particularly useful method for converting a compound of formula (I) to a different compound of formula (I) comprises reacting a compound of formula (I-G) (i.e., a compound of formula (I) wherein R^2 is fluoro) with an amine, and optionally heating the mixture to 50-150°C to prepare a compound of formula (I-H) (i.e., a compound of formula (I) wherein R^2 is $-NR^7R^8$).



wherein all variables are as defined above.

This procedure may be carried out by mixing a compound of formula (I-G) in an amine neat, or in a suitable solvent with an excess of amine to produce a compound of formula (I-H). Typically the solvent is a lower alcohol such as methanol, ethanol, isopropanol or the like. Other suitable solvents may include *N,N*-dimethylformamide, 1-methyl-2-pyrrolidine or the like.

As a further example, a compound of formula (I-I) may be converted to a compound of formula (I-J) using either of two methods.



wherein:

q is 0, 1 or 2;

Rg is Ay or Het;

M¹ is B(OH)₂, B(ORa)₂, B(Ra)₂, Sn(Ra)₃, Zn-halide, Zn-Ra or Mg-halide, where Ra is alkyl or cycloalkyl, halide is halo;

R^{1c} is Ay, NHR¹⁰Ay, -NR⁷Ay, Het, -NHHet, and -NHR¹⁰Het; and

all other variables are as defined above.

According to one process, a compound of formula (I-I) may be converted to a compound of formula (I-J) by a process comprising replacing the C-7 halogen of the compound of formula (I-I) with an amine. Typically the replacement is carried out by mixing the compound of formula (I-I) with an amine nucleophile of formula R^{1c} where R^{1c} is selected from the group consisting of -NR⁷R⁸, -NR⁷Ay, Het, -NHR¹⁰Het, NHHet, and -NHR¹⁰Ay; and optionally heating the reaction.

The reaction can also be carried out via an adaptation of procedures found in the literature (Wolfe, J. P.; Buchwald, S. L. *J. Org. Chem.* 2000, 65, 1144) wherein a compound of formula (I-I) is treated with an amine, a palladium (0) or nickel (0) source and a base in a suitable solvent. Suitable sources of palladium (0) include but
5 are not limited to palladium(II) acetate and tris(dibenzylideneacetone) dipalladium (0). Typical bases for use in the reaction include, for example sodium *tert*-butoxide and cesium carbonate. Toluene is an example of a suitable solvent.


Alternatively, a compound of formula (I-I) may be converted to a compound of
10 formula (I-J) by a process comprising coupling the compound of formula (I-I) with a metal compound of the formula $Rg-M^1$ where M^1 is $B(OH)_2$, $B(ORa)_2$, $B(Ra)_2$, $Sn(Ra)_3$, Zn-halide, Zn-Ra or Mg-halide, wherein Ra is alkyl or cycloalkyl and halide is halo. This process can be conveniently performed in an inert solvent, in the presence of a palladium (0) catalyst, optionally with heating. The reaction is performed by reacting
15 equimolar amounts of a compound of formula (I-I) with the metal compound of formula $Rg-M^1$ or optionally adding an excess of the metal compound. The palladium catalyst is preferably present in 1-10 mol% compared to the compound of formula (I-I). Palladium catalysts that may be used may include, but are not limited to, tetrakis(triphenylphosphine) palladium (0) dichlorobis(triphenylphosphine)palladium(II),
20 and bis(diphenylphosphinoferrocene)-palladium (II) dichloride. Inert solvents for use in the reaction include but are not limited to, *N,N*-dimethylformamide, toluene, tetrahydrofuran, dioxane, and 1-methyl-2-pyrrolidinone.

When the metal compound of formula $Rg-M^1$ is an arylboronic acid or ester or an
25 arylborinate, the reaction is more conveniently carried out by adding a base in a proportion equivalent to, or greater than, that of the metal compound.

Metal compounds of the formula $Rg-M^1$ can be purchased from commercial sources or prepared either as discreet isolated compounds or generated *in situ* by using methods
30 known to one skilled in the art. (Suzuki, A. *J. Organomet. Chem.* 1999, 576, 147; Stille,

J. Angew. Chem. Int. Ed. Engl. 1986, 25, 508; Snieckus, V. J. Org. Chem. 1995, 60, 292.)

As further example, a compound of formula (I-K) can be converted to a compound of formula (I-L) by reductive amination.



wherein:

R^{5a} is $-C(O)R^9$ or $-R^{10}C(O)R^9$;


R^{5b} is $-R^{10}Het$, $-R^{10}NR^7R^8$, $-R^{10}NR^7Ay$ or $-R^{10}NHHet$; and

all other variables are defined above.

By way of example this process for converting a compound of formula (I-K) to a compound of formula (I-L) can be performed by treating a compound of formula (I-K) with an amine in an inert solvent with a reducing agent, optionally in the presence of an acid. Numerous techniques for reductive amination are known to those skilled in the art and can be found in the literature. This conversion may be effected using any such conventional techniques and reagents. Typical conditions include but are not limited to solvents such as 1,2-dichloroethane, dichloromethane, tetrahydrofuran, ether, methanol and the like. Reducing agents by way of example include sodium borohydride, sodium triacetoxy borohydride and sodium cyanoborohydride. The ketone or aldehyde compound of formula (I-K) may be mixed with the amine prior to addition of reducing agent to prepare an intermediate imine *in situ*, followed by reduction to an amine of formula (I-L).

Another process for converting a compound of formula (I) to another compound of formula (I) involves by way of example treating a compound of formula (I-M) with an

amine nucleophile selected from the group consisting of Het-H, HNR^7R^8 , HNR^7Ay and NHHet , to prepare a compound of formula (I-N).



wherein:

X^1 is Cl, Br or iodo;


10 R^{5c} is $-\text{R}^{10}\text{Het}$, $-\text{R}^{10}\text{NR}^7\text{R}^8$, $-\text{R}^{10}\text{NR}^7\text{Ay}$ or $\text{R}^{10}\text{NHHet}$; and

all other variables are as defined above.

In particular, this method can be carried out by treating a compound of formula (I-M) with an amine nucleophile neat or in an inert solvent optionally with heating.

15 Preferred solvents include but are not limited to acetonitrile, dimethyl sulfoxide, tetrahydrofuran, *N,N*-dimethylformamide, dioxane, 1-methyl-2-pyrrolidinone, and the lower alcohols such as methanol, ethanol, isopropanol and the like.


20 In another example, a compound of formula (I-O) can be converted to a compound of formula (I-P) by a hydrolysis reaction.



This can be performed in a biphasic or miscible solvent system which includes an aqueous component under acidic conditions. Typical solvents include but are not limited to water, tetrahydrofuran, dioxane, methanol, acetone and the like.

30 Another example of conversion of a compound of formula (I) to another compound of formula (I) involves the conversion of a compound of formula (I-Q) to a compound of

formula (I-R) by oxidation. A compound of formula (I-Q) may be prepared by the process described in Scheme 1 above or by reaction of a compound of formula (I-P) with a suitable organo-metallic reagent, such as a Grignard reagent. A compound of formula (I-Q) may further be converted to a compound of formula (I-S) by reduction.



wherein:

R^d is selected from H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, -R¹⁰cycloalkyl, -R¹⁰Ay, -R¹⁰Het, -R¹⁰OR⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷Ay, -R¹⁰NHHet, and Ay; and

20 all other variables are defined above in connection with scheme 1.

By way of example the foregoing process of oxidizing a compound of formula (I-Q) to prepare a compound of formula (I-R) can be performed in an inert solvent in the presence of a conventional oxidant. Typical oxidants include but are not limited to manganese dioxide, and pyridinium chlorochromate and the like. Suitable solvents include but are not limited to dichloromethane, chloroform, tetrahydrofuran and the like.

In addition, a compound of formula (I-Q) can be reduced to prepare a compound of general formula (I-S) by treatment with a reducing agent in a suitable solvent. General techniques for reducing include but are not limited to catalytic

30

hydrogenation, and hydride reagents and the like. Preferred catalytic hydrogenation conditions typically include the use of palladium on carbon in approximately 10 weight percent of substrate and an atmosphere of hydrogen at ambient or elevated pressures. Suitable solvents include but are not limited to methanol, ethanol, ethyl acetate and the like. In addition, the reduction may be accomplished through the use of hydride reagents such as lithium aluminum hydride in an ethereal solvent such as tetrahydrofuran and the like. Yet another reduction technique suitable for the conversion of a compound of formula (I-Q) to a compound of formula (I-S) involves the use of triethylsilane in the presence of boron trifluoride etherate in an inert solvent such as tetrahydrofuran or dichloromethane, and the like.

Based upon this disclosure and the examples contained herein one skilled in the art can readily convert a compound of formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof into another compound of formula (I), or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

The present invention also provides radiolabeled compounds of formula (I) and biotinylated compounds of formula (I). Radiolabeled compounds of formula (I) and biotinylated compounds of formula (I) can be prepared using conventional techniques. For example, radiolabeled compounds of formula (I) can be prepared by reacting the compound of formula (I) with tritium gas in the presence of an appropriate catalyst to produce radiolabeled compounds of formula (I).


In one preferred embodiment, the compounds of formula (I) are tritiated.

The radiolabeled compounds of formula (I) and the biotinylated compounds of formula (I) are useful in assays for the identification of compounds for the treatment or prophylaxis of viral infections such as herpes viral infections. Accordingly, the present invention provides an assay method for identifying compounds which have activity for the treatment or prophylaxis of viral infections such as herpes viral

infections, which method comprises the step of specifically binding the radiolabeled compound of formula (I) or the biotinylated compounds of formula (I) to the target protein. More specifically, suitable assay methods will include competition binding assays. The radiolabeled compounds of formula (I) can be employed in assays
5 according to the methods conventional in the art.

The following examples are illustrative embodiments of the invention, not limiting the scope of the invention in any way, the invention being defined by the claims which follow. Reagents are commercially available or are prepared according to procedures
10 in the literature. Example numbers refer to those compounds listed in the tables above. ¹H and ¹³C NMR spectra were obtained on Varian Unity Plus NMR spectrophotometers at 300 or 400 MHz, and 75 or 100 MHz respectively. ¹⁹F NMR were recorded at 282 MHz. Mass spectra were obtained on Micromass Platform, or ZMD mass spectrometers from Micromass Ltd. Altrincham, UK, using either
15 Atmospheric Chemical Ionization (APCI) or Electrospray Ionization (ESI). Analytical thin layer chromatography was used to verify the purity of some intermediates which could not be isolated or which were too unstable for full characterization, and to follow the progress of reactions. Unless otherwise stated, this was done using silica gel (Merck Silica Gel 60 F254). Unless otherwise stated, column chromatography for
20 the purification of some compounds, used Merck Silica gel 60 (230–400 mesh), and the stated solvent system under pressure. All compounds were characterized as their free-base form unless otherwise stated. On occasion the corresponding hydrochloride salts were formed to generate solids where noted.

25 Example 1: 3-(2-Fluoropyridin-4-yl)-2-propylpyrazolo[1,5-a]pyridine.




a) 2-Fluoro-4-pent-1-ynylpyridine.

To a cold (0 °C) solution of 2-fluoro-4-iodopyridine (1.0 g, 4.5 mmol) in tetrahydrofuran (15 mL) was added copper (I) iodide (86 mg, 0.45 mmol), dichlorobis(triphenylphosphine) palladium (II) (157 mg, 0.22 mmol), and triethylamine (1.8 mL, 13.4 mmol). Pentyne (0.88 mL, 8.9 mmol) was added dropwise to the reaction mixture and the resultant solution was stirred at 0 °C for 30 minutes and then at room temperature for 18 hours. Water and ether were added and the layers separated. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (5:1 hexane-ether) provided 2-fluoro-4-pent-1-ynylpyridine (670 mg, 92%) as a clear oil. ¹H NMR (CDCl₃) δ 8.11 (d, 1 H), 7.11 (d, 1 H), 6.88 (s, 1 H), 2.41 (t, 2 H), 1.63 (m, 2 H), 1.04 (t, 3 H); ¹⁹F NMR (CDCl₃) δ -68.68; MS *m/z* 164 (M+1).

b) 3-(2-Fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine.


To a cold (0 °C) solution of 2-fluoro-4-pent-1-ynylpyridine (650 mg, 3.98 mmol) in acetonitrile (15 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (0.89 mL, 5.97 mmol) and 1-aminopyridinium iodide (974 mg, 4.38 mmol). The resultant dark solution was allowed to warm to room temperature and stirred vigorously. An additional equivalent of 1,8-diazabicyclo[5.4.0]undec-7-ene and 1-aminopyridinium iodide were added after 18 and again after 36 hours. Water was added and the mixture concentrated *in vacuo*. The residue was partitioned between water and ether. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (4:1 to 3:1 hexane-ethyl acetate) provided 3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine (765 mg, 75%) as an off-white solid. ¹H NMR (CDCl₃) δ 8.48 (d, 1 H), 8.30 (d, 1 H), 7.64 (d, 1 H), 7.32-7.23 (m, 2 H), 7.04 (s, 1 H), 6.86 (t, 1 H), 2.95 (t, 2 H), 1.82 (m, 2 H), 1.03 (t, 3 H); ¹⁹F NMR (CDCl₃) δ -68.48; MS *m/z* 256 (M+1).

Example 2: N-Cyclopentyl-4-(2-propylpyrazolo[1,5-a]pyridin-3-yl)pyridin-2-amine.



A solution of 3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-a]pyridine (126 mg, 0.49 mmol) in cyclopentylamine (5 mL) was heated in a sealed vessel at 150 °C for 18 h. The mixture was cooled to room temperature and concentrated *in vacuo* and the residue was purified by flash chromatography on silica gel (2:1 hexane-ethyl acetate) to give N-cyclopentyl-4-(2-propylpyrazolo[1,5-a]pyridin-3-yl)pyridin-2-amine (155 mg, 98%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.41 (d, 1 H), 8.07 (d, 1 H), 7.58 (d, 1 H), 7.14 (t, 1 H), 6.74 (td, 1 H), 6.66 (dd, 1 H), 6.45 (s, 1 H), 5.01 (broad, 1 H), 3.95 (m, 1 H), 2.91 (t, 2 H), 2.04 (m, 2 H), 1.80-1.50 (m, 8 H), 0.98 (t, 3 H); MS *m/z* 321 (M+1).

Example 3: 7-Chloro-3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-a]pyridine.




A solution of 3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-a]pyridine (109 mg, 0.42 mmol) in tetrahydrofuran (5 mL) was cooled to -78 °C. Butyllithium (0.4 mL, 1.6 M in hexane, 0.64 mmol) was added dropwise and the resultant solution was stirred at -78 °C for 45 minutes. Carbon tetrachloride (0.3 mL) was added and the mixture was stirred at -78 °C for 30 minutes and then quenched by the addition of methanol while cold. Upon warming to room temperature water and ether were added and the layers separated. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (4:1 to 2:1 hexane-ethyl acetate) provided 7-chloro-3-(2-fluoropyridin-4-yl)-2-

propylpyrazolo[1,5-*a*]pyridine (77 mg, 63%) as an orange solid. ^1H NMR (CDCl_3) δ 8.33 (d, 1 H), 7.58 (d, 1 H), 7.30 (m, 1 H), 7.22 (t, 1 H), 7.02–6.98 (m, 2 H), 2.99 (t, 2 H), 1.80 (m, 2 H), 1.01 (t, 3 H); ^{19}F NMR (CDCl_3) δ -68.11; MS m/z 290 ($M+1$).

5 Example 4: *N*-Cyclopentyl-3-[2-(cyclopentylamino)pyridin-4-yl]-2-propylpyrazolo[1,5-*a*]pyridin-7-amine.

10




A solution of 7-chloro-3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine (41 mg, 0.14 mmol) in cyclopentylamine (5 mL) was heated in a sealed vessel at 150 °C for 15 18 h. The mixture was cooled to room temperature and concentrated *in vacuo* and the residue was purified by flash chromatography on silica gel (2:1 hexane-ethyl acetate) to give *N*-cyclopentyl-3-[2-(cyclopentylamino)pyridin-4-yl]-2-propylpyrazolo[1,5-*a*]pyridin-7-amine (44 mg, 77%) as a yellow solid. ^1H NMR (CDCl_3) δ 8.10 (d, 1 H), 7.18 (t, 1 H), 7.00 (d, 1 H), 6.73 (d, 1 H), 6.52 (s, 1 H), 5.97–5.91 (m, 2 H), 5.07 (m, 1 H), 4.00 (m, 2 H), 2.95 (t, 2 H), 2.18–2.05 (m, 4 H), 1.87–1.57 (m, 14 H), 1.02 (t, 3 H); MS m/z 404 ($M+1$).

20

Example 5: 2-Isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine.

25




a) 4-(4-Methylpent-1-ynyl)-2-(methylthio)pyrimidine.

To a cold (0 °C) solution of 4-iodo-2-(methylthio)pyrimidine (4.73 g, 21.2 mmol) in tetrahydrofuran (70 mL) was added copper (I) iodide (404 mg, 0.212 mmol), dichlorobis(triphenylphosphine) palladium (II) (744 mg, 1.06 mmol), and triethylamine (8.8 mL, 63.8 mmol). 4-Methyl-pentyne (5.0 mL, 42.5 mmol) was added dropwise to the reaction mixture and the resultant solution was stirred at 0 °C for 30 minutes and then at room temperature for 72 hours. Water and ether were added and the layers separated. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (4:1 hexane-ether) provided 4-(4-methylpent-1-ynyl)-2-(methylthio)pyrimidine (4.0 g, 92%) as a clear oil. ¹H NMR (CDCl₃) δ 8.44 (d, 1 H), 6.96 (d, 1 H), 2.56 (s, 3 H), 2.35 (d, 2 H), 1.97 (m, 1 H), 1.05 (d, 6 H); MS *m/z* 207 (M+1).

b) 2-Isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine.


To a cold (0 °C) solution of 4-(4-methylpent-1-ynyl)-2-(methylthio)pyrimidine (2.5 g, 12 mmol) in acetonitrile (40 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (2.7 mL, 18 mmol) and 1-aminopyridinium iodide (3.49 g, 15.7 mmol). The resultant dark solution was allowed to warm to room temperature and stirred vigorously. The mixture was heated to 50 °C for 18 h then cooled to room temperature. Water was added and the mixture concentrated *in vacuo*. The residue was partitioned between water and ether. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (6:1 to 4:1 to 2:1 hexane-ethyl acetate) provided 2-isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine (2.0 g, 56%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.51-8.48 (m, 2 H), 8.36 (d, 1 H), 7.35 (t, 1 H), 7.19 (d, 1 H), 6.91 (t, 1 H), 3.04 (d, 2 H), 2.68 (s, 3 H), 2.20 (m, 1 H), 1.03 (d, 6 H); MS *m/z* 299 (M+1).

Example 6: 2-Isobutyl-3-[2-(methylsulfinyl)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine.



To a cold (0 °C) solution of 2-isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine (890 mg, 2.98 mmol) in dichloromethane (25 mL) was added sodium bicarbonate solid (500 mg), and *m*-chloroperbenzoic acid (734 mg, ~70% pure, 2.98 mmol). The mixture was stirred at 0 °C for 20 minutes and then quenched by the addition of saturated aqueous sodium thiosulfate and saturated aqueous sodium bicarbonate. The biphasic mixture was allowed to warm to room temperature and stirred vigorously for 15 minutes. The layers were separated and the organic layer washed with brine. The aqueous layer was extracted with dichloromethane and the combined organics were dried over sodium sulfate. Filtration and concentration *in vacuo* provided 2-isobutyl-3-[2-(methylsulfinyl)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine (930 mg, 99%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.73 (d, 1 H), 8.63 (d, 1 H), 8.47 (d, 1 H), 7.48 (d, 1 H), 7.43 (t, 1 H), 6.94 (t, 1 H), 3.00 (d, 2 H), 2.99 (s, 3 H), 2.16 (m, 1 H), 1.01 (d, 6 H) ; MS *m/z* 315 (M+1).

Example 7: *N*-Cyclopentyl-4-(2-isobutylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine.




A solution of 2-isobutyl-3-[2-(methylsulfinyl)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine (930 mg, 2.98 mmol) in cyclopentylamine (5 mL) was heated at 130 °C for 4 h. The mixture was cooled to room temperature and concentrated *in vacuo* and the residue was purified by flash chromatography on silica gel (2:1 to 1:1 hexane-ethyl acetate) to

give *N*-cyclopentyl-4-(2-isobutylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine (890 mg, 89%) as a White solid. ¹H NMR (CDCl₃) δ 8.44 (d, 1 H), 8.28-8.25 (m, 2 H), 7.26 (m, 1 H), 6.85-6.76 (m, 2 H), 5.08 (m, 1 H), 4.40 (m, 1 H), 3.00 (d, 2 H), 2.26-2.05 (m, 3 H), 1.83-1.51 (m, 6 H), 1.00 (d, 6 H); MS *m/z* 336 (M+1).

5

Example 8: *N*-Cyclopentyl-4-[2-isobutyl-7-(methylthio)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine.

10




15

To a cold (-78 °C) solution of *N*-cyclopentyl-4-(2-isobutylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine (750 mg, 2.23 mmol) in tetrahydrofuran (25 mL) was added lithium diisopropylamide (LDA) (11.18 mL of a 0.6 M stock solution was prepared from 18.75 mL of 1.6 M butyllithium and 4.20 mL of diisopropylamine in 27 mL of tetrahydrofuran, 6.71 mmol) dropwise. The resultant solution was stirred at -78 °C for 15 minutes. Dimethyldisulfide (1.0 mL, 11.15 mmol) was added rapidly and the mixture was stirred for 1 hour. Water and ether were added and the layers separated. The organic layer was washed with brine. The aqueous layer was extracted with ether and the combined organics were dried over magnesium sulfate. Filtration and concentration followed flash chromatography on silica gel (2:1 hexane-ether) provided *N*-cyclopentyl-4-[2-isobutyl-7-(methylthio)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine. (712 mg, 84%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.25 (d, 1 H), 8.07 (d, 1 H), 7.25 (t, 1 H), 6.75 (d, 1 H), 6.61 (d, 1 H), 5.04 (d, 1 H), 4.37 (m, 1 H), 3.05 (d, 2 H), 2.61 (s, 3 H), 2.26-2.05 (m, 3 H), 1.79-1.50 (m, 6 H), 0.97 (d, 6 H); MS *m/z* 382 (M+1).


25

Example 9: N-Cyclopentyl-4-[2-isobutyl-7-(methylsulfinyl)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine.




To a cold (0 °C) solution of N-cyclopentyl-4-[2-isobutyl-7-(methylthio)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine (700 mg, 1.83 mmol) in dichloromethane (30 mL) was added sodium bicarbonate solid (500 mg), and *m*-chloroperbenzoic acid (451 mg, ~70% pure, 1.83 mmol). The mixture was stirred at 0 °C for 20 minutes and then quenched by the addition of saturated aqueous sodium thiosulfate and saturated aqueous sodium bicarbonate. The biphasic mixture was allowed to warm to room temperature and stirred vigorously for 15 minutes. The layers were separated and the organic layer washed with brine. The aqueous layer was extracted with dichloromethane and the combined organics were dried over sodium sulfate. Filtration and concentration *in vacuo* provided N-cyclopentyl-4-[2-isobutyl-7-(methylsulfinyl)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine (720 mg, 99%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.38-8.29 (m, 2 H), 7.51-7.44 (m, 2 H), 6.77 (d, 1 H), 5.24 (m, 1 H), 4.38 (m, 1 H), 3.13 (s, 3 H), 3.08-2.91 (m, 2 H), 2.25-2.05 (m, 3 H), 1.82-1.54 (m, 6 H), 0.98 (m, 6 H); MS *m/z* 398 (M+1).

Example 10: N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isobutylpyrazolo[1,5-*a*]pyridin-7-amine.




A solution of *N*-cyclopentyl-4-[2-isobutyl-7-(methylsulfinyl)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine (110 mg, 0.28 mmol) in cyclopentylamine (5 mL) was heated at 130 °C for 72 h. The mixture was cooled to room temperature and concentrated *in vacuo* and the residue was purified by flash chromatography on silica gel (hexane-ethyl acetate) to give *N*-cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isobutylpyrazolo[1,5-*a*]pyridin-7-amine (93 mg, 81%) as a White solid. ¹H NMR (CDCl₃) δ 8.20 (d, 1 H), 7.50 (d, 1 H), 7.21 (t, 1 H), 6.74 (d, 1 H), 5.97-5.92 (m, 2 H), 5.09 (d, 1 H), 4.38 (m, 1 H), 3.96 (m, 1 H), 3.01 (d, 2 H), 2.27-2.04 (m, 5 H), 1.81-1.50 (m, 12 H), 0.97 (d, 6 H); MS *m/z* 419 (M+1).

Example 11: 2-(Diethoxymethyl)-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine.



The title compound was prepared in a similar manner to those described above to give a tinted oil. ¹H NMR (CDCl₃) δ 8.57-8.47 (m, 3 H), 7.79 (d, 1 H), 7.34 (t, 1 H), 6.92 (t, 1 H), 5.92 (s, 1 H), 3.80-3.62 *m, 4 H), 2.64 (s, 3 H), 1.23 (t, 6 H); MS *m/z* 345 (M+1).


Example 12: 3-[2-(Methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine-2-carbaldehyde.



A solution of 2-(diethoxymethyl)-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine (1.3 g, 3.8 mmol) in tetrahydrofuran (20 mL) was treated with aqueous hydrochloric acid (10 mL, 4 M). the resultant solution was stirred for 18 h. The solvents were removed *in vacuo* and the residue was taken up in dichloromethane and the pH adjusted to >7 with 1N aqueous sodium hydroxide. The organic layer was


washed with brine. The aqueous layer was extracted with dichloromethane and the combined organics were dried over sodium sulfate. Filtration and concentration followed flash chromatography on silica gel (1:1 to 1:2 hexane-ethyl acetate) provided 3-[2-(Methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine-2-carbaldehyde (804 mg, 79%) as a yellow solid. ¹H NMR (CDCl₃) δ 10.42 (s, 1 H), 8.61-8.58 (m, 3 H), 7.92 (d, 1 H), 7.44 (m, 1 H), 7.12 (m, 1 H), 2.65 (s, 3 H); MS *m/z* 271 (M+1).

Example 13: {3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol.




To a cold (-78 °C) solution of 3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine-2-carbaldehyde (520 mg, 1.92 mmol) in tetrahydrofuran (40 mL) was added phenylmagnesium bromide (2.88 mL, 1.0 M in tetrahydrofuran, 2.88 mmol). The reaction was warmed to 0 °C and stirred for 1 hour. The reaction mixture was then recooled to -78 °C and additional phenylmagnesium bromide (2.00 mL, 1.0 M in tetrahydrofuran, 2.00 mmol) was added and stirred at room temperature for 14 hours. The reaction was quenched with water and the resulting mixture was concentrated *in vacuo* to remove the excess tetrahydrofuran. The resulting aqueous mixture was extracted with dichloromethane. The organic layer was washed with water and dried over magnesium sulfate. Filtration and concentration, followed by flash chromatography (49:1 dichloromethane:methanol) provided {3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol (550 mg, 82%) as a white solid. R_f 0.31 (49:1 dichloromethane:methanol); ¹H NMR (CDCl₃) δ 8.52 (d, 1H), 8.40 (d, 1H), 7.87 (d, 1H), 7.43-7.13 (m, 7H), 6.93 (t, 1H), 6.21 (s, 1H), 2.60 (s, 3H); MS *m/z* 349 (M+1).

Example 14: {3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}{phenyl)methanol.



To a cold (0 °C) solution of {3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridin-2-yl}{phenyl)methanol (294 mg, 0.844 mmol) in dichloromethane (30 mL) was added *m*-chloroperoxybenzoic acid (218 mg, 1.26 mmol). The reaction mixture was stirred for 1.5 hours at 0 °C, then diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate solution. The organic layer was dried over magnesium sulfate, then filtered and concentrated. The resulting solid was heated in cyclopentylamine (2 mL, 20 mmol) at 45 °C for 16 hours. Concentration and flash chromatography (59:1 dichloromethane:methanol) provided {3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}{phenyl)methanol (120 mg, 89%) as a light yellow solid. *R*_f 0.55 (29:1 dichloromethane:methanol); ¹H NMR (CDCl₃) δ 8.50 (d, 1H), 8.22 (d, 1H), 7.96 (br, 1H), 7.82 (d, 1H), 7.40 (d, 2H), 7.35-7.17 (m, 4H), 6.85 (t, 1H), 6.78 (d, 1H), 6.20 (s, 1H), 5.12 (s, 1H), 4.29 (m, 1H), 2.05 (m, 2H), 1.79-1.60 (m, 4H), 1.49 (m, 2H); MS *m/z* 386 (M+1).


Example 15: {3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}{phenyl)methanone.



To a solution of {3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}{phenyl)methanol (227 mg, 0.589 mmol) in chloroform (10 mL) was added manganese dioxide (454 mg, 5.22 mmol). The reaction mixture was stirred for 2 days


at room temperature. The reaction mixture was diluted with dichloromethane and filtered through a pad of celite. The filtrate was concentrated to provide {3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanone (200 mg, 89%) as a white solid. R_f 0.30 (1:1 hexanes:ethyl acetate); ¹H NMR (CDCl₃) δ 8.58 (d, 1H), 8.40 (d, 1H), 8.16 (d, 1H), 8.00 (d, 2H), 7.61 (t, 1H), 7.55-7.38 (m, 3H), 7.01 (t, 1H), 6.75 (d, 1H), 5.02 (d, 1H), 4.18 (m, 1H), 1.96 (m, 2H), 1.70-1.41 (m, 6H); MS *m/z* 384 (M+1).

Example 16: {7-(Cyclopentylamino)-3-[2-(cyclopentylamino)-4-pyrimidinyl]-pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanone.




The title compound was prepared in a similar manner to those described above to give a Yellow solid; R_f 0.44 (3:2 hexanes:ethyl acetate); ¹H NMR (CDCl₃) δ 8.11 (d, 1H), 7.98 (d, 2H), 7.51 (d, 2H), 7.49-7.38 (m, 3H), 6.70 (d, 1H), 6.10 (d, 1H), 6.01 (d, 1H), 4.89 (d, 1H), 4.04-3.96 (m, 2H), 2.12 (m, 2H), 1.94-1.15 (m, 14H); MS *m/z* 467 (M+1).

Example 17: 4-(2-Benzylpyrazolo[1,5-*a*]pyridin-3-yl)-N-cyclopentyl-2-pyrimidinamine.



To a solution of concentrated sulfuric acid (9 mL, 169 mmol) in ethanol (25 mL) was added 10% palladium on carbon (200 mg) and a solution of {3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol (350 mg, 0.908 mmol) in ethanol (5 mL). The reaction mixture was stirred under 55 PSI of hydrogen for 3 days. The reaction mixture was filtered through a pad of celite. Saturated aqueous sodium bicarbonate solution was added to the filtrate until basic. The resulting mixture was concentrated on rotovap to remove the excess ethanol, then extracted with ethyl acetate. The organic layer was washed with water and brine, then dried over magnesium sulfate. Filtration and concentration, followed by flash chromatography (29:1 dichloromethane:methanol) provided 4-(2-benzylpyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine (221 mg, 66%) as an off-white solid. ¹H NMR (CDCl₃) δ 8.45 (d, 1H), 8.25 (d, 1H), 8.19 (d, 1H), 7.30-7.13 (m, 6H), 6.82 (m, 1H), 6.63 (d, 1H), 5.12 (d, 1H), 4.52 (s, 2H), 4.35 (m, 1H), 2.03 (m, 2H), 1.78-1.50 (m, 6H); MS *m/z* 370 (M+1).


Example 18: 4-(2-Benzyl-7-chloropyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine.



To a cold (-78 °C) solution of 4-(2-benzylpyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine (207 mg, 0.560 mmol) in tetrahydrofuran (7 mL) was added lithium diisopropylamide (17.5 mL, 0.16 M in tetrahydrofuran, 2.80 mmol). The reaction mixture was stirred at -78 °C for 45 minutes, then carbon tetrachloride (542 µL, 5.60 mmol) was added. The reaction mixture was stirred for an additional 30 minutes at -78 °C. The reaction mixture was quenched with water, warmed to room temperature, and extracted with ethyl acetate. The organic layer was washed with water and brine, then dried over magnesium sulfate. Filtration and concentration, followed by flash


chromatography (4:1 hexanes:ethyl acetate) provided 4-(2-benzyl-7-chloropyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine (60 mg, 27%) as a yellow solid. ¹H NMR (CDCl₃) δ 8.35 (d, 1H), 8.21 (d, 1H), 7.35-7.16 (m, 6H), 7.04 (d, 1H), 6.64 (d, 1H), 5.20 (d, 1H), 4.60 (s, 2H), 4.37 (m, 1H), 2.09 (m, 2H), 1.82-1.48 (m, 6H); MS *m/z* 404 (M+1).

Example 19: *N*-{4-[2-Benzyl-7-(cyclopentylamino)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinyl}-*N*-cyclopentylamine.



A solution of 4-(2-benzyl-7-chloropyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine (65 mg, 0.16 mmol) in cyclopentylamine (2 mL, 20 mmol) was heated at 116 °C for 16 hours. The reaction mixture was cooled and the excess cyclopentylamine was removed *in vacuo*. The crude material was chromatographed (19:1 dichloromethane:acetone) to provide *N*-{4-[2-benzyl-7-(cyclopentylamino)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinyl}-*N*-cyclopentylamine (23 mg, 32%) as a pale yellow solid. R_f 0.12 (5:1 hexanes:ethyl acetate); ¹H NMR δ 8.18 (d, 1H), 7.59 (d, 1H), 7.30-7.10 (m, 6H), 6.65 (d, 1H), 5.99 (d, 1H), 4.59 (s, 2H), 4.30 (m, 1H), 3.98 (m, 1H), 2.21-2.00 (m, 4H), 1.90-1.50 (m, 12H); MS *m/z* 453 (M+1).

Example 20: *N*-Cyclopentyl-4-[2-(methoxymethyl)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinamine.




The title compound was prepared in a similar manner to those described above to give a tan solid; R_f 0.32 (1:1 hexanes:ethyl acetate); ¹H NMR δ 8.49 (d, 1H), 8.40 (d, 1H), 8.30 (d, 1H), 7.30 (t, 1H), 6.99 (d, 1H), 6.90 (t, 1H), 5.21 (d, 1H), 4.90 (s, 2H), 4.39 (m, 1H), 3.50 (s, 3H), 2.12 (m, 2H), 1.82-1.50 (m, 6H); MS *m/z* 324 (M+1).

5

Example 21: N-Cyclopentyl-4-[2-(methoxymethyl)-7-(methylsulfanyl)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinamine.

10




The title compound was prepared in a similar manner to those described above to give a green solid; R_f 0.26 (1:1 hexanes:ethyl acetate); ¹H NMR (CDCl₃) δ 8.38-8.28 (m, 2H), 7.31 (m, 1H), 7.08 (d, 1H), 6.70 (d, 1H), 5.21 (d, 1H), 4.90 (s, 2H), 4.39 (m, 1H), 3.50 (s, 3H), 2.65 (s, 3H), 2.15 (m, 2H), 1.85-1.50 (m, 6H); MS *m/z* 370 (M+1).

15

Example 22: N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-(methoxymethyl)pyrazolo[1,5-*a*]pyridin-7-amine.

20



25


The title compound was prepared in a similar manner to those described above to give a yellow solid; R_f 0.21 (2:1 hexanes:ethyl acetate); ¹H NMR (CDCl₃) δ 8.23 (d, 1H), 7.65 (d, 1H), 7.29 (m, 1H), 6.05-5.95 (m, 2H), 5.15 (d, 1H), 4.89 (s, 2H), 4.40 (m, 1H), 3.99 (m, 1H), 3.51 (s, 3H), 2.20-2.02 (m, 4H), 1.90-1.50 (m, 12H); MS *m/z* 407 (M+1).

30

Example 23: N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(1-pyrrolidinyl)propyl]pyrazolo[1,5-a]pyridin-7-amine.

5

10




The title compound was prepared in a similar manner to those described above to give a brown solid; ^1H NMR (CDCl_3) δ 8.24 (d, 1H), 7.51 (d, 1H), 7.29 (m, 1H), 6.80 (d, 1H), 6.04–5.98 (m, 2H), 5.18 (d, 1H), 4.40 (m, 1H), 4.03 (m, 1H), 3.24 (t, 2H), 3.88–3.62 (br, 6H), 2.35–2.10 (m, 6H), 2.00–1.55 (m, 12); MS m/z 474 ($M+1$).

15


Example 24: N-({3-[2-(Methylsulfanyl)-4-pyrimidinyl]pyrazolo[1,5-a]pyridin-2-yl}methyl)-2-propanamine.

20




The title compound was prepared in a similar manner to those described above to give a yellow solid; ^1H NMR (CDCl_3) δ 8.52–8.48 (m, 2H), 8.16 (d, 1H), 7.41–7.34 (m, 2H), 6.95 (t, 1H), 4.35 (s, 2H), 3.07 (m, 1H), 2.62 (s, 3H), 1.25 (d, 6H); MS m/z 314 ($M+1$).

Example 25: *N*-Cyclopentyl-4-{2-[(isopropylamino)methyl]pyrazolo[1,5-*a*]pyridin-3-yl}-2-pyrimidinamine.




The title compound was prepared in a similar manner to those described above to give a beige solid; *R_f* 0.50 (47:3 dichloromethane:methanol); ¹H NMR (CDCl₃) δ 8.42 (d, 1H), 8.25 (d, 1H), 8.19 (d, 1H), 7.23 (m, 1H), 6.90 (d, 1H), 6.76 (t, 1H), 5.30 (br, 1H), 4.35 (m, 1H), 4.20 (s, 2H), 2.90 (m, 1H), 2.07 (m, 2H), 1.80–1.42 (m, 6H), 1.10 (d, 6H); MS *m/z* 351 (M+1).

Example 26: *N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(isopropylamino)methyl]pyrazolo[1,5-*a*]pyridin-7-amine.




The title compound was prepared in a similar manner to those described above to give a brown solid. ¹H NMR (CDCl₃) δ 11.30 (br, 1H), 8.40 (d, 1H), 7.41 (t, 1H), 7.25 (m, 2H), 6.89 (d, 1H), 6.15 (d, 1H), 6.02 (d, 1H), 4.52 (s, 2H), 4.36 (m, 1H), 4.02 (m, 1H), 2.99 (br, 1H), 2.22–1.95 (m, 4H), 1.95–1.55 (m, 12 H), 1.42 (d, 6H); MS *m/z* 434 (M+1).

Example 27: 4-{7-Chloro-2-[3-(isopropylamino)propyl]pyrazolo[1,5-*a*]pyridin-3-yl}-*N*-cyclopentyl-2-pyrimidinamine.




The title compound was prepared in a similar manner to those described above to give a tan solid. ¹H NMR (CDCl₃) δ 8.31 (d, 1H), 8.20 (d, 1H), 7.20 (t, 1H), 6.98 (d, 1H), 6.80 (d, 1H), 5.29 (br, 1H), 4.38 (m, 1H), 3.22 (t, 2H), 2.79 (m, 1H), 2.70 (t, 2H), 2.15-1.98 (m, 4H), 1.80-1.50 (m, 6H), 1.04 (d, 6H); MS *m/z* 413 (M+1).

Example 28: *N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(isopropylamino)propyl]pyrazolo[1,5-*a*]pyridin-7-amine.




The title compound was prepared in a similar manner to those described above to give a brown oil. ¹H NMR (CDCl₃) δ 8.26 (d, 1H), 7.50 (d, 1H), 7.30 (d, 1H), 6.80 (d, 1H), 6.02-5.95 (m, 2H), 5.20 (br, 1H), 4.40 (m, 1H), 4.01 (m, 1H), 3.22 (t, 2H), 2.80 (m, 1H), 2.70 (t, 2H), 2.21-1.98 (m, 6H), 1.98-1.50 (m, 12H), 1.07 (d, 6H); MS *m/z* 462 (M+1).

Example 29: 4-{7-Chloro-2-[(2-methoxyethoxy)methyl]pyrazolo[1,5-a]pyridin-3-yl}-N-cyclopentyl-2-pyrimidinamine.




The title compound was prepared in a similar manner to those described above to give a tan solid; ^1H NMR (CDCl_3) δ 8.50 (d, 1H), 8.35 (d, 1H), 7.27 (d, 1H), 7.20 (d, 1H), 7.03 (d, 1H), 5.21 (d, 1H), 4.98 (s, 2H), 4.38 (m, 1H), 3.78 (m, 2H), 3.58 (m, 2H), 3.37 (s, 3H), 2.09 (m, 2H), 1.80-1.49 (m, 6H); MS m/z 402 ($M+1$).

Example 30: 3-[2-(Cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)methyl]-N-(2-methoxyethyl)pyrazolo[1,5-a]pyridin-7-amine.




The title compound was prepared in a similar manner to those described above to give a yellow oil; ^1H NMR (CDCl_3) δ 8.27 (d, 1H), 7.76 (d, 1H), 7.30 (m, 1H), 7.10 (d, 1H), 6.32 (t, 1H), 6.02 (d, 1H), 5.18 (d, 1H), 5.00 (s, 2H), 4.41 (m, 1H), 3.81-3.70 (m, 4H), 3.62-3.50 (m, 4H), 3.46 (s, 3H), 3.41 (s, 3H), 2.15 (m, 2H), 1.83-1.52 (m, 6H); MS m/z 441 ($M+1$).

Example 31: N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)methyl]pyrazolo[1,5-a]pyridin-7-amine.



The title compound was prepared in a similar manner to those described above to give a yellow solid; ¹H NMR (CDCl₃) δ 8.29 (d, 1H), 7.64 (d, 1H), 7.30 (t, 1H), 7.09 (d, 1H), 6.09–5.99 (m, 2H), 5.20 (d, 1H), 5.00 (s, 2H), 4.41 (m, 1H), 4.00 (m, 1H), 3.80 (m, 2H), 3.62 (m, 2H), 3.41 (s, 3H), 2.22–2.02 (m, 4H), 1.95–1.50 (m, 12H); MS *m/z* 451 (M+1).


Example 32: N-Cyclopentyl-4-(2-isopropylpyrazolo[1,5-a]pyridin-3-yl)pyrimidin-2-amine.



The title compound was prepared in a similar manner as described above to give a brown solid. ¹H NMR (CDCl₃) δ: 8.46 (d, 1H), 8.28 (d, 1H), 8.15 (d, 1H), 7.23 (m, 1H), 6.84–6.76 (m, 2H), 5.08 (d, 1H), 4.38 (m, 1H), 2.14–2.04 (m, 4H), 1.89–1.53 (m, 4H), 1.45 (d, 6H); MS *m/z* 322 (M+1).

Example 33: N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isopropylpyrazolo[1,5-a]pyridin-7-amine.

5




The title compound was prepared in a similar manner as described above to give a brown solid. ^1H NMR (CDCl_3) δ : 8.16 (d, 1H), 7.41 (d, 1H), 7.21 (d, 1H), 6.76 (d, 1H), 6.00, (d, 1H), 5.93 (d, 1H), 4.37 (m, 1H), 3.95 (m, 1H), 2.134–2.04 (3H, m), 1.83–1.62 (m, 8H), 1.41 (d, 6H), 1.30–1.22 (m, 5H); MS m/z 405 ($M+1$).

10

Example 34: 2-Cyclopropyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-a]pyridine.

15




The title compound was prepared in a similar manner as described above to give a yellow solid. ^1H NMR (CDCl_3) δ : 8.50–8.39 (m, 3H), 7.54 (d, 1H), 7.30 (m, 1H), 6.87 (m, 1H), 2.67 (s, 3H), 2.31 (m, 1H), 1.16 (m, 4H); MS m/z 283 ($M+1$).

20

Example 35: N-Cyclopentyl-4-(2-cyclopropylpyrazolo[1,5-a]pyridin-3-yl)pyrimidin-2-amine.


25



The title compound was prepared in a similar manner as described above to give a tan solid. ^1H NMR (CDCl_3) δ : 8.36 (d, 2H), 8.27 (d, 1H), 7.23 (m, 1H), 7.10 (d, 1H), 6.79 (m,

1H), 5.06 (d, 1H), 4.38 (m, 1H), 2.36 (m, 1H), 2.13-2.07 (m, 2H), 1.80-1.53 (m, 6H), 1.18-1.06 (m, 4H); MS m/z 320 (M+1)

Example 36: N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-cyclopropylpyrazolo[1,5-a]pyridin-7-amine.



The title compound was prepared in a similar manner as described above to give a yellow solid. ¹H NMR (CDCl₃) δ: 8.22 (d, 1H), 7.66 (d, 1H), 7.23 (m, 1H), 7.08 (d, 1H), 5.93 (d, 1H), 5.87 (d, 1H), 5.09 (d, 1H), 4.39 (m, 1H), 3.95 (m, 1H), 2.43 (m, 1H), 2.14-2.07 (m, 4H), 1.88-1.50 (m, 12H), 1.15 (m, 2H), 1.07 (m, 2H); MS m/z 403 (M+1)

Example 37: Biological Activity

In the following example, "MEM" means Minimal Essential Media; "FBS" means Fetal Bovine Serum; "NP40" and "Igepal" are detergents; "MOI" means Multiplicity of Infection; "NaOH" means sodium hydroxide; "MgCl₂" means magnesium chloride; "dATP" means deoxyadenosine 5' triphosphate; "dUTP" means deoxyuridine 5' triphosphate; "dCTP" means dioxycytidine 5' triphosphate; "dGTP" means deoxyguanosine 5' triphosphate; "GuSCN" means Guanidinium thiocyanate; "EDTA" means ethylenediamine tetraacetic acid; "TE" means Tris-EDTA; "SCC" means sodium chloride/sodium citrate; "APE" means a solution of ammonia acetate, ammonia phosphate, EDTA; "PBS" means phosphate buffered saline; and "HRP" means horseradish peroxidase.

a) Tissue Culture and HSV infection.

Vero 76 cells were maintained in MEM with Earle's salts, L-glutamine, 8% FBS

(Hyclone, A-1111-L) and 100 units/mL Penicillin-100 µg/mL Streptomycin. For assay conditions, FBS was reduced to 2%. Cells are seeded into 96-well tissue culture plates

at a density of 5×10^4 cells/well after being incubated for 45 min at 37°C in the presence of HSV-1 or HSV-2 (MOI = 0.001). Test compounds are added to the wells and the plates are incubated at 37°C for 40–48 hours. Cell lysates are prepared as follows: media was removed and replaced with 150 µL/well 0.2 N NaOH with 1%
5 Igepal CA 630 or NP-40. Plates were incubated up to 14 days at room temperature in a humidified chamber to prevent evaporation.

(b) Preparation of detection DNA.

For the detection probe, a gel-purified, digoxigenin-labeled, 710-bp PCR fragment of
10 the HSV UL-15 sequence was utilized. PCR conditions included 0.5 µM primers, 180 µM dTTP, 20 µM dUTP-digoxigenin (Boehringer Mannheim 1558706), 200 µM each of dATP, dCTP, and dGTP, 1X PCR Buffer II (Perkin Elmer), 2.5 mM MgCl₂, 0.025 units/µL of AmpliTaq Gold polymerase (Perkin Elmer), and 5 ng of gel-purified HSV DNA per 100 µL. Extension conditions were 10 min at 95°C, followed by 30 cycles of 95°C for 1
15 min, 55°C for 30 sec, and 72°C for 2 min. The amplification was completed with a 10-min incubation at 72°C. Primers were selected to amplify a 728 bp probe spanning a section of the HSV1 UL15 open reading frame (nucleotides 249–977). Single-stranded transcripts were purified with Promega M13 Wizard kits. The final product was mixed 1:1 with a mixture of 6 M GuSCN, 100 mM EDTA and 200 µg/mL herring sperm DNA
20 and stored at 4°C.

(c) Preparation of capture plates.

The capture DNA plasmid (HSV UL13 region in pUC) was linearized by cutting with Xba I, denatured for 15 min at 95°C and diluted immediately into Reacti-Bind DNA
25 Coating Solution (Pierce, 17250, diluted 1:1 with TE buffer, pH 8) at 1 ng/µL. 75 µL/well were added to Corning (#3922 or 9690) white 96-well plates and incubated at room temperature for at least 4 hrs before washing twice with 300 µL/well 0.2X SSC/0.05% Tween-20 (SSC/T buffer). The plates were then incubated overnight at room temperature with 150 µL/well 0.2 N NaOH, 1% IGEPAL and 10 µg/mL herring
30 sperm DNA.

(d) Hybridization.

Twenty-seven (27) μL of cell lysate was combined with 45 μL of hybridization solution (final concentration: 3M GuSCN, 50 mM EDTA, 100 $\mu\text{g/ml}$ salmon sperm DNA, 5X Denhardt's solution, 0.25X APE, and 5 ng of the digoxigenin-labeled detection probe).

- 5 APE is 1.5 M NH_4 -acetate, 0.15 M ammonium phosphate monobasic, and 5 mM EDTA adjusted to pH 6.0. Mineral oil (50 μL) was added to prevent evaporation. The hybridization plates were incubated at 95°C for 10 minutes to denature the DNA, then incubated at 42°C overnight. The wells were washed 6X with 300 $\mu\text{L/well}$ SSC/T buffer then incubated with 75 $\mu\text{L/well}$ anti-digoxigenin- HRP-conjugated antibody
- 10 (Boehringer Mannheim 1207733, 1:5000 in TE) for 30 min at room temperature. The wells were washed 6X with 300 $\mu\text{L/well}$ with PBS/0.05% Tween-20 before 75 $\mu\text{L/well}$ SuperSignal LBA substrate (Pierce) was added. The plates were incubated at room temperature for 30 minutes and chemiluminescence was measured in a Wallac Victor reader.

15

e) Results.

The following results were obtained for HSV-1.

Example No.	IC ₅₀ (μM)
1	>40
2	8
3	37
4	0.9
5	21
6	24
7	18
8	3.8
9	1.9
10	0.5
12	>5
14	2.5
15	7


Example No.	IC ₅₀ (μM)
16	0.4
17	2
19	0.8
20	15
21	4.6
22	1.2
23	>2.5
24	2.1
25	27
26	>2.5
27	23
28	>5
29	23
30	4.7
31	1.5
32	4.4
33	2
34	>40
35	3.1
36	0.9

The results demonstrate that the compounds of the present invention are useful for the treatment and prophylaxis of herpes viral infections.

CLAIMS

1. A compound of formula (I):

5



wherein:

p is 0, 1, 2, 3 or 4;

- 10 each R¹ is the same or different and is independently selected from the group consisting of halo, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Ay, Het, -OR⁷, -OAY, -OR¹⁰AY, -OHet, -OR¹⁰Het, -C(O)R⁹, -C(O)AY, -C(O)Het, -CO₂R⁹, -C(O)NR⁷R⁸, -C(O)NR⁷AY, -C(O)NHR¹⁰AY, -C(O)NHR¹⁰Het, -C(S)NR⁹R¹¹, -C(NH)NR⁷R⁸, -C(NH)NR⁷AY, -S(O)_nR⁹, -S(O)_nAY, -S(O)_nHet, -S(O)₂NR⁷R⁸, -S(O)₂NR⁷AY, -NR⁷R⁸, -NR⁷AY, -NHHet, -NHR¹⁰AY, -NHR¹⁰Het, -R¹⁰cycloalkyl, -R¹⁰AY, -R¹⁰Het, -R¹⁰O-C(O)R⁹, -R¹⁰O-C(O)AY, -R¹⁰O-C(O)Het, -R¹⁰O-S(O)_nR⁹, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹, -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(O)NR⁷AY, -R¹⁰C(O)NHR¹⁰Het, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO₂R⁹, -R¹⁰SO₂NR⁹R¹¹, -R¹⁰SO₂NHCO⁹, -R¹⁰NR⁷R⁸, -R¹⁰NR⁷AY, -R¹⁰NHC(NH)NR⁹R¹¹, cyano, nitro and azido; or
- 20 two adjacent R¹ groups together with the atoms to which they are bonded form a C₅₋₆cycloalkyl or a 5 or 6-membered heterocyclic ring containing 1 or 2 heteroatoms;

- each R⁷ and R⁸ are the same or different and are independently selected from
- 25 the group consisting of H, alkyl, alkenyl, cycloalkyl, cycloalkenyl, -C(O)R⁹, -CO₂R⁹, -C(O)NR⁹R¹¹, -C(S)NR⁹R¹¹, -C(NH)NR⁹R¹¹, -SO₂R¹⁰, -SO₂NR⁹R¹¹, -R¹⁰cycloalkyl, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰CO₂R⁹, -R¹⁰C(O)NR⁹R¹¹, -R¹⁰C(S)NR⁹R¹¹, -R¹⁰C(NH)NR⁹R¹¹, -R¹⁰SO₂R¹⁰, -R¹⁰SO₂NR⁹R¹¹, -R¹⁰SO₂NHCO⁹, -R¹⁰NR⁹R¹¹, -R¹⁰NHCO⁹, -R¹⁰NHSO₂R⁹ and -R¹⁰NHC(NH)NR⁹R¹¹;
- 30

each R^9 and R^{11} are the same or different and are independently selected from the group consisting of H, alkyl, cycloalkyl, $-R^{10}$ cycloalkyl, $-R^{10}OH$, $-R^{10}(OR^{10})_w$ where w is 1-10, and $-R^{10}NR^{10}R^{10}$;

each R^{10} is the same or different and is independently selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkenyl;

Ay is aryl;

Het is a 5- or 6-membered heterocyclic or heteroaryl group;

R^2 is selected from the group consisting of halo, alkyl, alkenyl, cycloalkyl, cycloalkenyl,

Ay, Het, $-OR^7$, $-OAY$, $-OHet$, $-OR^{10}Het$, $-S(O)_nR^9$, $-S(O)_nAy$, $-S(O)_nNR^7R^8$,

$-S(O)_nHet$, $-NR^7R^8$, $-NHHet$, $-NHR^{10}Ay$, $-NHR^{10}Het$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$;

n is 0, 1 or 2;

Y is N or CH;

R^3 and R^4 are the same or different and are each independently selected from the

group consisting of H, halo, alkyl, alkenyl, cycloalkyl, Ay, Het, $-OR^7$, $-OAY$,

$-C(O)R^7$, $-C(O)AY$, $-CO_2R^7$, $-CO_2AY$, $-SO_2NHR^9$, $-NR^7R^8$, $-NR^7Ay$, $-NHHet$,

$-NHR^{10}Het$, $-R^{10}$ cycloalkyl, $-R^{10}OR^7$, $-R^{10}OAY$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$;

R^5 is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl,

cycloalkyl, cycloalkenyl, $-OR^7$, $-OAY$, $-OHet$, $-OR^{10}Ay$, $-OR^{10}Het$, $-C(O)R^9$,

$-C(O)AY$, $-C(O)Het$, $-CO_2R^9$, $-C(O)NR^7R^8$, $-C(O)NR^7Ay$, $-C(O)NHR^{10}Het$, $-CH(OR^9)_2$,

$-CH(OR^9)-R^{10}$, $-CH(OR^9)-AY$, $-C(S)NR^9R^{11}$, $-C(NH)NR^7R^8$, $-C(NH)NR^7Ay$, $-S(O)_nR^9$,

$-S(O)_2NR^7R^8$, $-S(O)_2NR^7Ay$, $-NR^7R^8$, $-NR^7Ay$, $-NHHet$, $-NHR^{10}Ay$, $-NHR^{10}Het$,

$-R^{10}$ cycloalkyl, $-R^{10}AY$, $-R^{10}Het$, $-R^{10}OR^9$, $-R^{10}C(O)R^9$, $-R^{10}C(O)AY$, $-R^{10}C(O)Het$,

$-R^{10}CO_2R^9$, $-R^{10}C(O)NR^9R^{11}$, $-R^{10}C(O)NR^7Ay$, $-R^{10}C(O)NHR^{10}Het$, $-R^{10}CH(OR^9)-R^{10}$;

$-R^{10}CH(OR^9)-AY$, $-R^{10}C(S)NR^9R^{11}$, $-R^{10}C(NH)NR^9R^{11}$, $-R^{10}SO_nR^9$, $-R^{10}SO_2NR^9R^{11}$,

$-R^{10}SO_2NHCOR^9$, $-R^{10}NR^7R^8$, $-R^{10}NR^7Ay$, $-R^{10}NHC(NH)NR^9R^{11}$, cyano, nitro and

azido; or

wherein when Y is CH, R^3 is not $-NR^7Ay$;

or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

2. The compound according to claim 1 wherein each R^1 is the same or different and is independently selected from the group consisting of halo, alkyl, cycloalkyl, Ay, Het, $-OR^7$, $-C(O)R^9$, $-C(O)Het$, $-CO_2R^9$, $-C(O)NR^7R^8$, $-C(O)NR^7Ay$, $-C(O)NHR^{10}Het$, $-S(O)_nR^9$, $-S(O)_2NR^7R^8$, $-S(O)_2NR^7Ay$, $-NR^7R^8$, $-NR^7Ay$, $-NHHet$, $-NHR^{10}Ay$, $-NHR^{10}Het$,
5 $-R^{10}cycloalkyl$, $-R^{10}Het$, $-R^{10}OR^9$, $-R^{10}C(O)NR^7Ay$, $-R^{10}SO_2NHCOR^9$, $-R^{10}NR^7R^8$, $-R^{10}NR^7Ay$, cyano, nitro and azido.
3. The compound according to claim 1 wherein each R^1 is the same or different and is independently selected from the group consisting of halo, Ay, Het, $-NR^7R^8$ and
10 $-NR^7Ay$.
4. The compound according to any of claims 1-3 wherein p is 0 or 1.
5. The compound according to any of claims 1-4 wherein R^2 is selected from the
15 group consisting of halo, alkenyl, cycloalkyl, cycloalkenyl, Ay, Het, $-OR^7$, $-OAY$, $-OHet$, $-OR^{10}Het$, $-S(O)_nR^9$, $-NR^7R^8$, $-NHHet$, $-NHR^{10}Het$, $-R^{10}NR^7R^8$ and $-R^{10}NR^7Ay$.
6. The compound according to any of claims 1-4 wherein R^2 is $-NR^7R^8$.
- 20 7. The compound according to any of claims 1-6 wherein Y is N.
8. The compound according to any of claims 1-6 wherein Y is CH.
9. The compound according to any of claims 1-8 wherein R^3 and R^4 are the same
25 or different and are each independently selected from the group consisting of H, halo, alkyl, Ay, $-OR^7$, $-CO_2R^7$, $-NR^7R^8$, $-R^{10}OR^7$ and $-R^{10}NR^7R^8$.
10. The compound according to any of claims 1-9 wherein R^3 and R^4 are both H.
- 30 11. The compound according to any of claims 1-10 wherein R^5 is selected from the group consisting of halo, alkyl, cycloalkyl, $-OR^7$, $-C(O)R^9$, $-C(O)Ay$, $-C(O)Het$,

-CH(OR⁹)-R¹⁰, -CH(OR⁹)-Ay, -S(O)_nR⁹, -S(O)₂NR⁷R⁸, -NR⁷R⁸, -NR⁷Ay, -R¹⁰cycloalkyl, -R¹⁰Ay, -R¹⁰Het, -R¹⁰OR⁹, -R¹⁰C(O)R⁹, -R¹⁰SO₂NR⁹R¹¹ and -R¹⁰NR⁷R⁸.

12. The compound according to any of claims 1-10, wherein R⁵ is selected from the group consisting of alkyl, -C(O)Ay, -CH(OR⁹)-Ay, -R¹⁰cycloalkyl, -R¹⁰Ay, -R¹⁰OR⁹ and -R¹⁰NR⁷R⁸.

13. A compound selected from the group consisting of:

3-(2-Fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine;

10 *N*-Cyclopentyl-4-(2-propylpyrazolo[1,5-*a*]pyridin-3-yl)pyridin-2-amine;

7-Chloro-3-(2-fluoropyridin-4-yl)-2-propylpyrazolo[1,5-*a*]pyridine;

N-Cyclopentyl-3-[2-(cyclopentylamino)pyridin-4-yl]-2-propylpyrazolo[1,5-*a*]pyridin-7-amine;

2-Isobutyl-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;

15 2-Isobutyl-3-[2-(methylsulfinyl)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;

N-Cyclopentyl-4-(2-isobutylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine;

N-Cyclopentyl-4-[2-isobutyl-7-(methylthio)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine;

N-Cyclopentyl-4-[2-isobutyl-7-(methylsulfinyl)pyrazolo[1,5-*a*]pyridin-3-yl]pyrimidin-2-amine;

20 2-amine;

N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-isobutylpyrazolo[1,5-*a*]pyridin-7-amine;

2-(Diethoxymethyl)-3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine;

3-[2-(Methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridine-2-carbaldehyde;

25 {3-[2-(methylthio)pyrimidin-4-yl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol;

{3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanol;

{3-[2-(Cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanone;

30 {7-(Cyclopentylamino)-3-[2-(cyclopentylamino)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}(phenyl)methanone;

- 4-(2-Benzylpyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine;
4-(2-Benzyl-7-chloropyrazolo[1,5-*a*]pyridin-3-yl)-*N*-cyclopentyl-2-pyrimidinamine;
N-{4-[2-Benzyl-7-(cyclopentylamino)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinyl}-*N*-
cyclopentylamine;
5 *N*-Cyclopentyl-4-[2-(methoxymethyl)pyrazolo[1,5-*a*]pyridin-3-yl]-2-pyrimidinamine;
N-Cyclopentyl-4-[2-(methoxymethyl)-7-(methylsulfanyl)pyrazolo[1,5-*a*]pyridin-3-yl]-
2-pyrimidinamine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-
(methoxymethyl)pyrazolo[1,5-*a*]pyridin-7-amine;
10 *N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(1-pyrrolidinyl)propyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
N-{3-[2-(Methylsulfanyl)-4-pyrimidinyl]pyrazolo[1,5-*a*]pyridin-2-yl}methyl-2-
propanamine;
N-Cyclopentyl-4-{2-[(isopropylamino)methyl]pyrazolo[1,5-*a*]pyridin-3-yl}-2-
15 pyrimidinamine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(isopropylamino)methyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
4-{7-Chloro-2-[3-(isopropylamino)propyl]pyrazolo[1,5-*a*]pyridin-3-yl}-*N*-
cyclopentyl-2-pyrimidinamine;
20 *N*-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[3-(isopropylamino)propyl]-
pyrazolo[1,5-*a*]pyridin-7-amine;
4-{7-Chloro-2-[(2-methoxyethoxy)methyl]pyrazolo[1,5-*a*]pyridin-3-yl}-*N*-
cyclopentyl-2-pyrimidinamine;
3-[2-(Cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)methyl]-*N*-(2-
25 methoxyethyl)pyrazolo[1,5-*a*]pyridin-7-amine;
N-Cyclopentyl-3-[2-(cyclopentylamino)-4-pyrimidinyl]-2-[(2-methoxyethoxy)-
methyl]pyrazolo[1,5-*a*]pyridin-7-amine;
N-Cyclopentyl-4-(2-isopropylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine;
N-Cyclo

N-Cyclopentyl-4-(2-cyclopropylpyrazolo[1,5-*a*]pyridin-3-yl)pyrimidin-2-amine; and
N-Cyclopentyl-3-[2-(cyclopentylamino)pyrimidin-4-yl]-2-cyclopropylpyrazolo[1,5-*a*]pyridin-7-amine;

or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

14. A pharmaceutical composition comprising a compound according to any of claims 1-13.

15. The pharmaceutical composition according to claim 14 further comprising a pharmaceutically acceptable carrier or diluent.

16. The pharmaceutical composition according to any of claims 14 or 15, further comprising an antiviral agent selected from the group consisting of aciclovir and valaciclovir or a pharmaceutically acceptable salt thereof.

17. A method for the prophylaxis or treatment of a herpes viral infection in an animal, said method comprising administering to the animal a therapeutically effective amount of a compound according to any of claims 1-13.

18. The method according to claim 17 wherein said herpes viral infection is selected from the group consisting of herpes simplex virus 1, herpes simplex virus 2, cytomegalovirus, Epstein Barr virus, varicella zoster virus, human herpes virus 6, human herpes virus 7, and human herpes virus 8.

19. A method for the prophylaxis or treatment of a condition or disease associated with a herpes viral infection in an animal, comprising administering to the animal a therapeutically effective amount of a compound according to any of claims 1-13.


20. A process for preparing a compound according to any of claims 1-13 comprising the steps of:

a) coupling a compound of formula (II):



wherein X is chloro, bromo, iodo or triflate;

10 to a terminal alkyne of formula (III):



to prepare a compound of formula (IV):



and

b) reacting an *N*-amino pyridinium salt of formula (V):



wherein Z⁻ is a counterion;

25 with the compound of the formula (IV) to prepare a compound of formula (I).

21. The process according to claim 20 further comprising the step of converting the compound of formula (I) to a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

30

22. The process according to claim 20 further comprising the step of converting the compound of formula (I) or a pharmaceutically acceptable salt, solvate or

physiologically functional derivative thereof to another compound of formula (I) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

- 5 23. A compound according to any of Claims 1-13 for use in therapy.
24. A compound according to any of Claims 1-13, for use in the prophylaxis or treatment of a herpes viral infection in an animal.
- 10 25. A compound according to any of Claims 1-13, for use in the prophylaxis or treatment of a condition or disease associated with a herpes viral infection in an animal.
- 15 26. The use of a compound according to any of claims 1-13 for the preparation of a medicament for the prophylaxis or treatment of a herpes viral infection in an animal.
- 20 27. The use of a compound according to any of claims 1-13 for the preparation of a medicament for the prophylaxis or treatment of a condition or disease associated with a herpes viral infection in an animal.
28. A pharmaceutical composition comprising a compound according to any of claims 1-13 for use in the prophylaxis or treatment of a herpes viral infection in an animal.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 03/30334

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D471/04 A61K31/437 A61P31/22 //(C07D471/04,231:00,221:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data, PAJ, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01/014375 A (BEATTIE JOHN FRANKLIN ; BREAULT GLORIA ANNE (GB); JEWSBURY PHILLIP JOH) 1 March 2001 (2001-03-01) claims 1,10,11	1-5,7, 14,15,23
A	WO 02/072581 A (SMITHKLINE BEECHAM CORP ; BOYD F LESLIE (US); JOHNS BRIAN A (US); GUDM) 19 September 2002 (2002-09-19) page 1, line 5 - line 8 page 4, formula I Examples	1-28
A	WO 02/48147 A (GLAXO GROUP LTD ; CHAMBERLAIN STANLEY D (US); JOHNS BRIAN A (US); GUDM) 20 June 2002 (2002-06-20) page 1, line 5 - line 8 page 5, formula I Examples	1-28
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 13 February 2004		Date of mailing of the international search report 23/02/2004
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer Hoepfner, W

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 03/30334

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 02/48148 A (GLAXO GROUP LTD ; CHAMBERLAIN STANLEY D (US); HARRIS PHILIP ANTHONY (U) 20 June 2002 (2002-06-20) page 1, line 5 - line 8 page 4, formula (I) Examples</p> <p>-----</p>	1-28

INTERNATIONAL SEARCH REPORT

Internat application No.
PCT/US 03/30334

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 17-19, 23-25 and 28 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Publication No
PCT/US 03/30334

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0114375	A	01-03-2001	AT 251623 T	15-10-2003
			AU 757639 B2	27-02-2003
			AU 6583300 A	19-03-2001
			BG 106383 A	30-09-2002
			BR 0013476 A	30-04-2002
			CA 2376293 A1	01-03-2001
			CN 1370163 T	18-09-2002
			CZ 20020617 A3	12-06-2002
			DE 60005850 D1	13-11-2003
			EE 200200080 A	16-06-2003
			EP 1214318 A1	19-06-2002
			WO 0114375 A1	01-03-2001
			HU 0202494 A2	28-10-2002
			JP 2003507478 T	25-02-2003
			NO 20020832 A	12-04-2002
			SK 2402002 A3	10-09-2002
			ZA 200200028 A	02-04-2003
WO 02072581	A	19-09-2002	EP 1366048 A2	03-12-2003
			WO 02072581 A2	19-09-2002
WO 0248147	A	20-06-2002	AU 3934402 A	24-06-2002
			AU 3934802 A	24-06-2002
			EP 1341788 A2	10-09-2003
			EP 1377573 A2	07-01-2004
			WO 0248147 A2	20-06-2002
			WO 0248148 A2	20-06-2002
WO 0248148	A	20-06-2002	AU 3934402 A	24-06-2002
			AU 3934802 A	24-06-2002
			EP 1341788 A2	10-09-2003
			EP 1377573 A2	07-01-2004
			WO 0248147 A2	20-06-2002
			WO 0248148 A2	20-06-2002